

Generators of KMS Symmetric Markov Semigroups on $\mathcal{B}(\mathfrak{h})$ Symmetry and Quantum Detailed Balance

F. FAGNOLA

Dipartimento di Matematica, Politecnico di Milano,
Piazza Leonardo da Vinci 32, I-20133 Milano (Italy)
franco.fagnola@polimi.it

V. UMANITÀ

Dipartimento di Matematica, Università di Genova,
Via Dodecaneso 35, I-16146 Genova (Italy)
veronica.umanita@fastwebnet.it

Abstract

We find the structure of generators of norm continuous quantum Markov semigroups on $\mathcal{B}(\mathfrak{h})$ that are symmetric with respect to the scalar product $\text{tr}(\rho^{1/2}x^*\rho^{1/2}y)$ induced by a faithful normal invariant state ρ and satisfy two quantum generalisations of the classical detailed balance condition related with this non-commutative notion of symmetry: the so-called standard detailed balance condition and the standard detailed balance condition with an antiunitary time reversal.

1 Introduction

Symmetric Markov semigroups have been extensively studied in classical stochastic analysis (Fukushima et al. [13] and the references therein) because their generators and associated Dirichlet forms are very well tractable by Hilbert space and probabilistic methods.

Their non-commutative counterpart has also been deeply investigated (Albeverio and Goswami [1], Cipriani [6], Davies and Lindsay [8], Goldstein and Lindsay [15], Guido, Isola and Scarlatti [17], Park [23], Sauvageot [26] and the references therein).

The classical notion of symmetry with respect to a measure, however, admits several non-commutative generalisations. Here we shall consider the so-called KMS-symmetry that seems more natural from a mathematical point of view (see e.g. Accardi and Mohari [3], Cipriani [6], [7], Goldstein and Lindsay [14], Petz [25]) and find the structure of generators of norm-continuous quantum Markov semigroups (QMS) on the von Neumann algebra $\mathcal{B}(\mathfrak{h})$ of all bounded operators on a complex separable Hilbert space $\mathfrak{h}$ that are symmetric or satisfy quantum detailed balance conditions associated with KMS-symmetry or generalising it.

We consider QMS on $\mathcal{B}(\mathfrak{h})$, i.e. weak*-continuous semigroups of normal, completely positive, identity preserving maps $\mathcal{T} = (\mathcal{T}_t)_{t \geq 0}$ on $\mathcal{B}(\mathfrak{h})$, with a faithful normal invariant state ρ . This defines pre-scalar products on $\mathcal{B}(\mathfrak{h})$ by $(x, y)_s = \text{tr}(\rho^{1-s} x^* \rho^s y)$ for $s \in [0, 1]$ and allows one to define the s -dual semigroup $\mathcal{T}'$ on $\mathcal{B}(\mathfrak{h})$ satisfying $\text{tr}(\rho^{1-s} x^* \rho^s \mathcal{T}_t(y)) = \text{tr}(\rho^{1-s} \mathcal{T}'_t(x)^* \rho^s y)$ for all $x, y \in \mathcal{B}(\mathfrak{h})$. The above scalar products coincide on an Abelian von Neumann algebra, the notion of symmetry $\mathcal{T} = \mathcal{T}'$, however, clearly depends on the choice of the parameter s .

The most studied cases are $s = 0$ and $s = 1/2$. Denoting $\mathcal{T}_*$ the predual semigroup, a simple computation yields $\mathcal{T}'(x) = \rho^{-(1-s)} \mathcal{T}_{*t}(\rho^{1-s} x \rho^s) \rho^{-s}$, and shows that for $s = 1/2$ the maps $\mathcal{T}'_t$ are positive but, for $s \neq 1/2$ this may not be the case. Indeed, it is well-known that, for $s \neq 1/2$, the maps $\mathcal{T}'_t$ are positive if and only if the maps $\mathcal{T}_t$ commute with the modular group $(\sigma_t)_{t \in \mathbb{R}}$, $\sigma_t(x) = \rho^{it} x \rho^{-it}$ (see e.g. [18] Prop. 2.1 p. 98, [22] Th. 6 p. 7985, for $s = 0$, [11] Th. 3.1 p. 341, Prop. 8.1 p. 362 for $s \neq 1/2$). This quite restrictive condition implies that the generator has a very special form that makes simpler the mathematical study of symmetry but imposes strong structural constraints (see e.g. [18] and [12]).

Here we shall consider the most natural choice $s = 1/2$ whose consequences are not so stringent and say that $\mathcal{T}$ is *KMS-symmetric* if it coincides with its dual $\mathcal{T}'$. KMS-symmetric QMS were introduced by Cipriani [6] and Goldstein and Lindsay [14]; we refer to [7] for a discussion of the connection with the KMS condition justifying this terminology.

All quantum versions of the classical principle of detailed balance (Agarwal [4], Alicki [5], Frigerio, Gorini, Kossakowski and Verri [18], Majewski [20], [21]), which is at the basis of equilibrium physics, are formulated prescribing a certain relationship between $\mathcal{T}$ and $\mathcal{T}'$ or between their generators, therefore they depending of the underlying notion of symmetry. This work clarifies the structure of generators of QMS that are KMS symmetric or satisfy a quantum detailed balance condition involving the above scalar product with $s = 1/2$ and is a key step towards understanding which is the most natural and flexible in view of the study of their generalisations for quantum

systems out of equilibrium as, for instance, the *dynamical* detailed balance condition introduced by Accardi and Imafuku [2].

The generator $\mathcal{L}$ of a norm-continuous QMS can be written in the standard Gorini-Kossakowski-Sudarshan [16] and Lindblad [19] (GKSL) form

$$\mathcal{L}(x) = i[H, x] - \frac{1}{2} \sum_{\ell \geq 1} (L_\ell^* L_\ell x - 2L_\ell^* x L_\ell + x L_\ell^* L_\ell) \quad (1)$$

where $H, L_\ell \in \mathcal{B}(\mathfrak{h})$ with $H = H^*$ and the series $\sum_{\ell \geq 1} L_\ell^* L_\ell$ is strongly convergent. The operators L_ℓ, H in (1) are not uniquely determined by $\mathcal{L}$, however, under a natural minimality condition (Theorem 8 below) and a zero-mean condition $\text{tr}(\rho L_\ell) = 0$ for all $\ell \geq 1$, H is determined up to a scalar multiple of the identity operator and the $(L_\ell)_{\ell \geq 1}$ up to a unitary transformation of the multiplicity space of the completely positive part of $\mathcal{L}$. We shall call *special* a GKSL representation of $\mathcal{L}$ by operators H, L_ℓ satisfying these conditions.

As a result, by the remark following Theorem 8, in a special GKSL representation of $\mathcal{L}$, the operator $G = -2^{-1} \sum_{\ell \geq 1} L_\ell^* L_\ell - iH$, is uniquely determined by $\mathcal{L}$ up to a purely imaginary multiple of the identity operator and allows us to write $\mathcal{L}$ in the form

$$\mathcal{L}(x) = G^* x + \sum_{\ell \geq 1} L_\ell^* x L_\ell + x G. \quad (2)$$

Our characterisation of QMS that are KMS-symmetric or satisfy a quantum detailed balance condition generalising related with KMS-symmetry is given in terms of the operators G, L_ℓ (or, in an equivalent way H, L_ℓ) of a special GKSL representation.

Theorem 18 shows that a QMS is KMS-symmetric if and only if the operators G, L_ℓ of a special GKSL representation of its generator satisfy $\rho^{1/2} G^* = G \rho^{1/2} + ic \rho^{1/2}$ for some $c \in \mathbb{R}$ and $\rho^{1/2} L_k^* = \sum_\ell u_{k\ell} L_\ell \rho^{1/2}$ for all k and some unitary $(u_{k\ell})$ on the multiplicity space of the completely positive part of $\mathcal{L}$ coinciding with its transpose, i.e. such that $u_{k\ell} = u_{\ell k}$ for all k, ℓ .

In order to describe our results on the structure of generators of QMS satisfying a quantum detailed balance condition we first recall some basic definitions. The best known is due to Alicki [5] and Frigerio-Gorini-Kossakowski-Verri [18]: a norm-continuous QMS $\mathcal{T} = (\mathcal{T}_t)_{t \geq 0}$ on $\mathcal{B}(\mathfrak{h})$ satisfies the *Quantum Detailed Balance* (QDB) condition if there exists an operator $\tilde{\mathcal{L}}$ on $\mathcal{B}(\mathfrak{h})$ and self-adjoint operator K on $\mathfrak{h}$ such that $\text{tr}(\rho \tilde{\mathcal{L}}(x)y) = \text{tr}(\rho x \mathcal{L}(y))$ and $\mathcal{L}(x) - \tilde{\mathcal{L}}(x) = 2i[K, x]$ for all $x, y \in \mathcal{B}(\mathfrak{h})$. Roughly speaking we can say that $\mathcal{L}$ satisfies the QDB condition if the difference of $\mathcal{L}$ and its adjoint $\tilde{\mathcal{L}}$ with respect to the pre-scalar product on $\mathcal{B}(\mathfrak{h})$ given by $\text{tr}(\rho a^* b)$ is a derivation.

This QDB implies that the operator $\tilde{\mathcal{L}} = \mathcal{L} - 2i[K, \cdot]$ is conditionally completely positive and then generates a QMS $\tilde{\mathcal{T}}$. Therefore $\mathcal{L}$ and the maps $\mathcal{T}_t$ commute with the modular group. This restriction does not follow if the dual QMS is defined with respect to the symmetric pre-scalar product with $s = 1/2$.

The QDB can be readily reformulated replacing $\tilde{\mathcal{L}}$ with the adjoint $\mathcal{L}'$ defined via the symmetric scalar product; the resulting condition will be called *Standard Quantum Detailed Balance* condition (SQDB) (see e.g. [9]).

Theorem 15 characterises generators $\mathcal{L}$ satisfying the SQDB and extends previous partial results by Park [23] and the authors [11]: the SQDB holds if and only if there exists a unitary matrix $(u_{k\ell})$, coinciding with its transpose, i.e. $u_{k\ell} = u_{\ell k}$ for all k, ℓ , such that $\rho^{1/2} L_k^* = \sum_{\ell} u_{k\ell} L_{\ell} \rho^{1/2}$. This shows, in particular, that the SQDB depends only on the L_{ℓ} 's and does not involve directly H and G . Moreover, we find explicitly the unitary $(u_{k\ell})_{k\ell}$ providing also a geometrical characterisation of the SQDB (Theorem 16) in terms of the operators $L_{\ell} \rho^{1/2}$ and their adjoints as Hilbert-Schmidt operators on $\mathfrak{h}$.

We also consider (Definition 5) another notion of quantum detailed balance, inspired by the original Agarwal's notion (see [4], Majewski [20], [21], Talkner [27]) involving an antiunitary *time reversal* operator θ which does not play any role in Alicki et al. definition. Time reversal appears to keep into the account the parity of quantum observables; position and energy, for instance, are even, i.e. invariant under time reversal, momentum are odd, i.e. change sign under time reversal. The original Agarwal's definition, however, depends on the $s = 0$ pre-scalar product and implies then, that a QMS satisfying this quantum detailed balance condition must commute with the modular automorphism. Here we study the modified version (Definition 5) involving the symmetric $s = 1/2$ pre-scalar product that we call the SQDB- θ condition.

Theorem 21 shows that $\mathcal{L}$ satisfies the SQDB- θ condition if and only if there exists a special GKSL representation of $\mathcal{L}$ by means of operators H, L_{ℓ} such that $G \rho^{1/2} = \rho^{1/2} \theta G^* \theta$ and a unitary self-adjoint $(u_{k\ell})_{k\ell}$ such that $\rho^{1/2} L_k^* = \sum_{\ell} u_{k\ell} \theta L_{\ell} \theta \rho^{1/2}$ for all k . Here again $(u_{k\ell})_{k\ell}$ is explicitly determined by the operators $L_{\ell} \rho^{1/2}$ (Theorem 22).

We think that these results show that the SQDB condition is somewhat weaker than the SQDB- θ condition because the first does not involve the directly the operators H, G . Moreover, the unitary operator in the linear relationship between $L_{\ell} \rho^{1/2}$ and their adjoints is transpose symmetric and any point of the unit disk could be in its spectrum while, for generators satisfying the SQDB- θ , it is self-adjoint and its spectrum is contained in $\{-1, 1\}$. Therefore, by the spectral theorem, it is possible in principle to find

a standard form for the generators of QMSs satisfying the SQDB- θ generalising the standard form of generators satisfying the usual QDB condition (that commute with the modular group) as illustrated in the case of QMSs on $M_2(\mathbb{C})$ studied in the last section. This classification must be much more complex for generators of QMSs satisfying the SQDB.

The above arguments and the fact that the SQDB- θ condition can be formulated in a simple way both on the QMS or on its generator (this is not the case for the QDB- θ when $\mathcal{L}$ and its Hamiltonian part $i[H, \cdot]$ do not commute), lead us to the conclusion that the SQDB- θ is the more natural non-commutative version of the classical detailed balance condition.

The paper is organised as follows. In Section 2 we construct the dual QMS $\mathcal{T}'$ and recall the quantum detailed balance conditions we investigate, then we study the relationship between the generators of a QMS and its adjoint in Section 3. Our main results on the structure of generators are proved in Sections 4 (QDB without time reversal) and 5 (with time reversal).

2 The dual QMS, KMS-symmetry and quantum detailed balance

We start this section by constructing the dual semigroup of a norm-continuous QMS with respect to the $(\cdot, \cdot)_{1/2}$ pre-scalar product on $\mathcal{B}(\mathfrak{h})$ defined by an invariant state ρ and prove some properties that will be useful in the sequel. Although this result may be known, the presentation given here leads in a simple and direct way to the dual QMS avoiding non-commutative L^p -spaces techniques.

Proposition 1 *Let Φ be a positive unital normal map on $\mathcal{B}(\mathfrak{h})$ with a faithful normal invariant state ρ . There exists a unique positive unital normal map Φ' on $\mathcal{B}(\mathfrak{h})$ such that*

$$\mathrm{tr}(\rho^{1/2}\Phi'(x)\rho^{1/2}y) = \mathrm{tr}(\rho^{1/2}x\rho^{1/2}\Phi(y))$$

for all $x, y \in \mathcal{B}(\mathfrak{h})$. If Φ is completely positive, then Φ' is also completely positive.

Proof. Let Φ_* be the predual map on the Banach space of trace class operators on $\mathfrak{h}$ and let $Rk(\rho^{1/2})$ denote the range of the operator $\rho^{1/2}$. This is clearly dense in $\mathfrak{h}$ because ρ is faithful and coincides with the domain of the unbounded self-adjoint operator $\rho^{-1/2}$.

For all self-adjoint $x \in \mathcal{B}(\mathfrak{h})$ consider the sesquilinear form on the domain $Rk(\rho^{1/2}) \times Rk(\rho^{1/2})$

$$F(v, u) = \langle \rho^{-1/2}v, \Phi_*(\rho^{1/2}x\rho^{1/2})\rho^{-1/2}u \rangle.$$

By the invariance of ρ and positivity of Φ_* we have

$$-\|x\|\rho = -\|x\|\Phi_*(\rho) \leq \Phi_*(\rho^{1/2}x\rho^{1/2}) \leq \|x\|\Phi_*(\rho) = \|x\|\rho.$$

Therefore $|F(u, u)| \leq \|x\| \cdot \|v\| \cdot \|u\|$. Thus sesquilinear form is bounded and there exists a unique bounded operator y such that, for all $u, v \in Rk(\rho^{1/2})$,

$$\langle v, yu \rangle = \langle \rho^{-1/2}v, \Phi_*(\rho^{1/2}x\rho^{1/2})\rho^{-1/2}u \rangle.$$

Note that, Φ being a $*$ -map, and x self/adjoint

$$\begin{aligned} \langle v, y^*u \rangle &= \overline{\langle y^*u, v \rangle} \\ &= \overline{\langle \rho^{-1/2}u, \Phi_*(\rho^{1/2}x\rho^{1/2})\rho^{-1/2}v \rangle} \\ &= \overline{\langle \Phi_*(\rho^{1/2}x\rho^{1/2})\rho^{-1/2}u, \rho^{-1/2}v \rangle} \\ &= \langle \rho^{-1/2}v, \Phi_*(\rho^{1/2}x\rho^{1/2})\rho^{-1/2}u \rangle. \end{aligned}$$

This shows that y is self-adjoint. Defining $\Phi'(x) := y$, we find a real-linear map on self-adjoint operators on $\mathcal{B}(\mathfrak{h})$ that can be extended to a linear map on $\mathcal{B}(\mathfrak{h})$ decomposing each self-adjoint operator as the sum of its self-adjoint and anti self-adjoint parts.

Clearly Φ' is positive because $\rho^{1/2}\Phi'(x^*x)\rho^{1/2} = \Phi_*(\rho^{1/2}x^*x\rho^{1/2})$ and Φ_* is positive. Moreover, by the above construction $\Phi'(\mathbb{1}) = \mathbb{1}$, i.e. Φ' is unital. Therefore Φ' is a norm-one contraction.

If Φ is completely positive, then Φ_* is also and formula $\rho^{1/2}\Phi'(x)\rho^{1/2} = \Phi_*(\rho^{1/2}x\rho^{1/2})$ shows that Φ' is completely positive.

Finally we show that Φ' is normal. Let $(x_\alpha)_\alpha$ be a net of positive operators on $\mathcal{B}(\mathfrak{h})$ with least upper bound $x \in \mathcal{B}(\mathfrak{h})$. For all $u \in \mathfrak{h}$ we have then

$$\begin{aligned} \sup_\alpha \langle \rho^{1/2}u, \Phi'(x_\alpha)\rho^{1/2}u \rangle &= \sup_\alpha \langle u, \Phi_*(\rho^{1/2}x_\alpha\rho^{1/2})u \rangle \\ &= \langle u, \Phi_*(\rho^{1/2}x\rho^{1/2})u \rangle = \langle \rho^{1/2}u, \Phi'(x)\rho^{1/2}u \rangle. \end{aligned}$$

Now if $u \in \mathfrak{h}$, for every $\varepsilon > 0$, we can find a $u_\varepsilon \in Rk(\rho^{1/2})$ such that $\|u - u_\varepsilon\| < \varepsilon$ by the density of the range of $\rho^{1/2}$. We have then

$$\begin{aligned} |\langle u, (\Phi'(x_\alpha) - \Phi'(x))u \rangle| &\leq \varepsilon \|\Phi'(x_\alpha) - \Phi'(x)\| (\|u\| + \|u_\varepsilon\|) \\ &\quad + |\langle u_\varepsilon, (\Phi'(x_\alpha) - \Phi'(x))u_\varepsilon \rangle| \end{aligned}$$

for all α . The conclusion follows from the arbitrariness of ε and the uniform boundedness of $\|\Phi'(x_\alpha) - \Phi'(x)\|$ and $\|u_\varepsilon\|$. $\square$

Theorem 2 *Let $\mathcal{T}$ be a QMS on $\mathcal{B}(\mathfrak{h})$ with a faithful normal invariant state ρ . There exists a QMS $\mathcal{T}'$ on $\mathcal{B}(\mathfrak{h})$ such that*

$$\rho^{1/2}\mathcal{T}'_t(x)\rho^{1/2} = \mathcal{T}_{*t}(\rho^{1/2}x\rho^{1/2}) \quad (3)$$

for all $x \in \mathcal{B}(\mathfrak{h})$ and all $t \geq 0$.

Proof. By Proposition 1, for each $t \geq 0$, there exists a unique completely positive normal and unital contraction $\mathcal{T}'_t$ on $\mathcal{B}(\mathfrak{h})$ satisfying (3). The semigroup property follows from the algebraic computation

$$\begin{aligned} \rho^{1/2}\mathcal{T}'_{t+s}(x)\rho^{1/2} &= \mathcal{T}_{*t}(\mathcal{T}_{*s}(\rho^{1/2}x\rho^{1/2})) \\ &= \mathcal{T}_{*t}(\rho^{1/2}\mathcal{T}'_s(x)\rho^{1/2}) = \rho^{1/2}\mathcal{T}'_t(\mathcal{T}'_s(x))\rho^{1/2}. \end{aligned}$$

Since the map $t \rightarrow \langle \rho^{1/2}v, \mathcal{T}'_t(x)\rho^{1/2}u \rangle$ is continuous by the identity (3) for all $u, v \in \mathfrak{h}$, and $\|\mathcal{T}'_t(x)\| \leq \|x\|$ for all $t \geq 0$, a 2ε approximation argument shows that $t \rightarrow \mathcal{T}'_t(x)$ is continuous for the weak*-operator topology on $\mathcal{B}(\mathfrak{h})$. It follows that $\mathcal{T}' = (\mathcal{T}'_t)_{t \geq 0}$ is a QMS on $\mathcal{B}(\mathfrak{h})$. $\square$

Definition 3 *The quantum Markov semigroup $\mathcal{T}'$ is called the dual semigroup of $\mathcal{T}$ with respect to the invariant state ρ .*

It is easy to see, using (3), that ρ is an invariant state also for $\mathcal{T}'$.

Remark 1 When $\mathcal{T}$ is norm-continuous it is not clear whether also $\mathcal{T}'$ is norm-continuous. Here, however, we are interested in generators of symmetric or detailed balance QMS. We shall see that these additional properties of $\mathcal{T}$ imply that also $\mathcal{T}'$ is norm continuous. Therefore we proceed studying norm-continuous QMSs whose dual is also norm-continuous.

The quantum detailed balance condition of Alicki, Frigerio, Gorini, Kosakowski and Verri modified by considering the pre-scalar product $(\cdot, \cdot)_{1/2}$ on $\mathcal{B}(\mathfrak{h})$, usually called *standard* (see e.g. [9]) because of multiplications by $\rho^{1/2}$ as in the standard representation of $\mathcal{B}(\mathfrak{h})$, is defined as follows.

Definition 4 *The QMS $\mathcal{T}$ generated by $\mathcal{L}$ satisfies the standard quantum detailed balance condition (SQDB) if there exists an operator $\mathcal{L}'$ on $\mathcal{B}(\mathfrak{h})$ and a self-adjoint operator K on $\mathfrak{h}$ such that*

$$\mathrm{tr}(\rho^{1/2}x\rho^{1/2}\mathcal{L}(y)) = \mathrm{tr}(\rho^{1/2}\mathcal{L}'(x)\rho^{1/2}y), \quad \mathcal{L}(x) - \mathcal{L}'(x) = 2i[K, x] \quad (4)$$

for all $x \in \mathcal{B}(\mathfrak{h})$.

The operator $\mathcal{L}'$ in the above definition must be norm-bounded because it is everywhere defined and norm closed. To see this consider a sequence $(x_n)_{n \geq 1}$ in $\mathcal{B}(\mathfrak{h})$ converging in norm to a $x \in \mathcal{B}(\mathfrak{h})$ such that $(\mathcal{L}(x_n))_{n \geq 1}$ converges in norm to $b \in \mathcal{B}(\mathfrak{h})$ and note that

$$\begin{aligned} \mathrm{tr}(\rho^{1/2} \mathcal{L}'(x) \rho^{1/2} y) &= \lim_{n \rightarrow \infty} \mathrm{tr}(\rho^{1/2} x_n \rho^{1/2} \mathcal{L}(y)) \\ &= \lim_{n \rightarrow \infty} \mathrm{tr}(\rho^{1/2} \mathcal{L}'(x_n) \rho^{1/2} y) = \mathrm{tr}(\rho^{1/2} b \rho^{1/2} y) \end{aligned}$$

for all $y \in \mathcal{B}(\mathfrak{h})$. The elements $\rho^{1/2} y \rho^{1/2}$, with $y \in \mathcal{B}(\mathfrak{h})$, are dense in the Banach space of trace class operators on $\mathfrak{h}$ because ρ is faithful. Therefore shows that $\mathcal{L}'(x) = b$ and $\mathcal{L}'$ is closed.

Since both $\mathcal{L}$ and $\mathcal{L}'$ are bounded, also K is bounded.

We now introduce another definition of quantum detailed balance, due to Agarwal [4] with the $s = 0$ pre-scalar product, that involves a *time reversal* θ . This is an antiunitary operator on $\mathfrak{h}$, i.e. $\langle \theta u, \theta v \rangle = \langle v, u \rangle$ for all $u, v \in \mathfrak{h}$, such that $\theta^2 = \mathbb{1}$ and $\theta^{-1} = \theta^* = \theta$.

Recall that, θ is antilinear, i.e. $\theta z u = \bar{z} u$ for all $u \in \mathfrak{h}$, $z \in \mathbb{C}$, and its adjoint θ^* satisfies $\langle u, \theta v \rangle = \langle v, \theta^* u \rangle$ for all $u, v \in \mathfrak{h}$. Moreover $\theta x \theta$ belongs to $\mathcal{B}(\mathfrak{h})$ (linearity is re-established) and $\mathrm{tr}(\theta x \theta) = \mathrm{tr}(x^*)$ for every trace-class operator x ([10] Prop. 4), indeed, taking an orthonormal basis of $\mathfrak{h}$, we have

$$\begin{aligned} \mathrm{tr}(\theta x \theta) &= \sum_j \langle e_j, \theta x \theta e_j \rangle = \sum_j \langle x \theta e_j, \theta^* e_j \rangle \\ &= \sum_j \langle \theta e_j, x^* \theta^* e_j \rangle = \mathrm{tr}(x^*). \end{aligned}$$

It is worth noticing that the cyclic property of the trace does not hold for θ , since $\mathrm{tr}(\theta x \theta) = \mathrm{tr}(x^*)$ may not be equal to $\mathrm{tr}(x)$ for non self-adjoint x .

Definition 5 *The QMS $\mathcal{T}$ generated by $\mathcal{L}$ satisfies the standard quantum detailed balance condition with respect to the time reversal θ (SQDB- θ) if*

$$\mathrm{tr}(\rho^{1/2} x \rho^{1/2} \mathcal{L}(y)) = \mathrm{tr}(\rho^{1/2} \theta y^* \theta \rho^{1/2} \mathcal{L}(\theta x^* \theta)), \quad (5)$$

for all $x, y \in \mathcal{B}(\mathfrak{h})$.

The operator θ is used to keep into the account parity of the observables under time reversal. Indeed, a self-adjoint operator $x \in \mathcal{B}(\mathfrak{h})$ is called *even* (resp. *odd*) if $\theta x \theta = x$ (resp. $\theta x \theta = -x$). The typical example of antilinear time reversal is a conjugation (with respect to some orthonormal basis).

This condition is usually stated ([20], [21], [27]) for the QMS $\mathcal{T}$ as

$$\mathrm{tr}(\rho^{1/2}x\rho^{1/2}\mathcal{T}_t(y)) = \mathrm{tr}(\rho^{1/2}\theta y^*\theta\rho^{1/2}\mathcal{T}_t(\theta x^*\theta)), \quad (6)$$

for all $t \geq 0$, $x, y \in \mathcal{B}(\mathfrak{h})$. In particular, for $t = 0$ we find that this identity holds if and only if ρ and θ commute, i.e. ρ is an even observable. This is the case, for instance, when ρ is a function of the energy.

Lemma 6 *The following conditions are equivalent:*

(i) θ and ρ commute,

(ii) $\mathrm{tr}(\rho^{1/2}x\rho^{1/2}y) = \mathrm{tr}(\rho^{1/2}\theta y^*\theta\rho^{1/2}\theta x^*\theta)$ for all $x, y \in \mathcal{B}(\mathfrak{h})$.

Proof. If ρ and θ commute, from $\mathrm{tr}(\theta a \theta) = \mathrm{tr}(a^*)$, we have

$$\mathrm{tr}(\rho^{1/2}\theta y^*\theta\rho^{1/2}\theta x^*\theta) = \mathrm{tr}(\theta(\rho^{1/2}\theta y^*\theta\rho^{1/2}x^*)\theta) = \mathrm{tr}(x\rho^{1/2}y\rho^{1/2})$$

and (ii) follows cycling $\rho^{1/2}$. Conversely, if (ii) holds, taking $x = \mathbb{1}$, we have

$$\mathrm{tr}(\rho y) = \mathrm{tr}(\rho\theta y^*\theta) = \mathrm{tr}(\theta(\theta y^*\theta)^*\rho\theta) = \mathrm{tr}(y\theta\rho\theta) = \mathrm{tr}(\theta\rho\theta y),$$

for all $y \in \mathcal{B}(\mathfrak{h})$, and $\rho = \theta\rho\theta$. □

Proposition 7 *If ρ and θ commute then (5) and (6) are equivalent.*

Proof. Clearly (5) follows from (6) differentiating at $t = 0$.

Conversely, putting $\alpha(x) = \theta x \theta$ and denoting $\mathcal{L}_*$ the predual of $\mathcal{L}$ we can write (5) as

$$\mathrm{tr}(\mathcal{L}_*(\rho^{1/2}x\rho^{1/2})y) = \mathrm{tr}(\rho^{1/2}\alpha(y^*)\rho^{1/2}\mathcal{L}(\alpha(x^*))) = \mathrm{tr}(\rho^{1/2}\alpha(\mathcal{L}(\alpha(x)))\rho^{1/2}y),$$

for all $y \in \mathcal{B}(\mathfrak{h})$, because $\mathrm{tr}(\alpha(a)) = \mathrm{tr}(a^*)$. Therefore we have

$$\mathcal{L}_*(\rho^{1/2}x\rho^{1/2}) = \rho^{1/2}\alpha(\mathcal{L}(\alpha(x)))\rho^{1/2}$$

and, iterating, $\mathcal{L}_*^n(\rho^{1/2}x\rho^{1/2}) = \rho^{1/2}\alpha(\mathcal{L}^n(\alpha(x)))\rho^{1/2}$ for all $n \geq 1$. It follows that (5) holds for all powers $\mathcal{L}^n$ with $n \geq 1$. Since ρ and θ commute, it is true also for $n = 0$ and we find (6) by the exponentiation formula $\mathcal{T}_t = \sum_{n \geq 0} t^n \mathcal{L}^n / n!$. □

We do not know whether the SQDB condition (4) of Definition 4 has a simple explicit formulation in terms of the maps $\mathcal{T}_t$ if $\mathcal{L}$ and $\mathcal{L}'$ do not commute.

Remark 2 The SQDB condition (5), by $\text{tr}(\theta a \theta) = \text{tr}(a^*)$, reads

$$\text{tr}(\rho^{1/2} x \rho^{1/2} \mathcal{L}(y)) = \text{tr}(\rho^{1/2} (\theta \mathcal{L}(\theta x \theta) \theta) \rho^{1/2} x),$$

for all $x, y \in \mathcal{B}(\mathfrak{h})$, i.e. $\mathcal{L}'(x) = \theta \mathcal{L}(\theta x \theta) \theta$.

Write $\mathcal{L}$ in a special GKSL form as in (1) and decompose the generator $\mathcal{L} = \mathcal{L}_0 + i[H, \cdot]$ into the sum of its dissipative part $\mathcal{L}_0$ and derivation part $i[H, \cdot]$. If H commutes with θ , by the antilinearity of θ , we find $\mathcal{L}'(x) = \theta \mathcal{L}_0(\theta x \theta) \theta - i[H, x]$. Therefore, if the dissipative part is time reversal invariant, i.e. $\mathcal{L}_0(x) = \theta \mathcal{L}_0(\theta x \theta) \theta$, we end up with $\mathcal{L}' = \mathcal{L} - 2i[H, \cdot]$.

The relationship with Definition 4 of SQDB, in this case, is then clear. The SQDB conditions of Definition 4 and 5, however, in general are not comparable.

3 The generator of a QMS and its dual

We shall always consider *special* GKSL representations of the generator of a norm-continuous QMS by means of operators L_ℓ, H . These are described by the following theorem (we refer to [24] Theorem 30.16 for the proof).

Theorem 8 *Let $\mathcal{L}$ be the generator of a norm-continuous QMS on $\mathcal{B}(\mathfrak{h})$ and let ρ be a normal state on $\mathcal{B}(\mathfrak{h})$. There exists a bounded self-adjoint operator H and a finite or infinite sequence $(L_\ell)_{\ell \geq 1}$ of elements of $\mathcal{B}(\mathfrak{h})$ such that:*

- (i) $\text{tr}(\rho L_\ell) = 0$ for each $\ell \geq 1$,
- (ii) $\sum_{\ell \geq 1} L_\ell^* L_\ell$ is a strongly convergent sum,
- (iii) if $\sum_{\ell \geq 0} |c_\ell|^2 < \infty$ and $c_0 + \sum_{\ell \geq 1} c_\ell L_\ell = 0$ for complex scalars $(c_k)_{k \geq 0}$ then $c_k = 0$ for every $k \geq 0$,
- (iv) the GKSL representation (1) holds.

If $H', (L'_\ell)_{\ell \geq 1}$ is another family of bounded operators in $\mathcal{B}(\mathfrak{h})$ with H' self-adjoint and the sequence $(L'_\ell)_{\ell \geq 1}$ is finite or infinite then the conditions (i)–(iv) are fulfilled with $H, (L_\ell)_{\ell \geq 1}$ replaced by $H', (L'_\ell)_{\ell \geq 1}$ respectively if and only if the lengths of the sequences $(L_\ell)_{\ell \geq 1}, (L'_\ell)_{\ell \geq 1}$ are equal and for some scalar $c \in \mathbb{R}$ and a unitary matrix $(u_{\ell j})_{\ell, j}$ we have

$$H' = H + c, \quad L'_\ell = \sum_j u_{\ell j} L_j.$$

As an immediate consequence of the uniqueness (up to a scalar) of the Hamiltonian H , the decomposition of $\mathcal{L}$ as the sum of the derivation $i[H, \cdot]$ and a dissipative part $\mathcal{L}_0 = \mathcal{L} - i[H, \cdot]$ determined by special GKSL representations of $\mathcal{L}$ is unique. Moreover, since $(u_{\ell j})$ is unitary, we have

$$\sum_{\ell \geq 1} (L'_\ell)^* L'_\ell = \sum_{\ell, k, j \geq 1} \bar{u}_{\ell k} u_{\ell j} L_k^* L_j = \sum_{k, j \geq 1} \left(\sum_{\ell \geq 1} \bar{u}_{\ell k} u_{\ell j} \right) L_k^* L_j = \sum_{k \geq 1} L_k^* L_k.$$

Therefore, putting $G = -2^{-1} \sum_{\ell \geq 1} L_\ell^* L_\ell - iH$, we can write $\mathcal{L}$ in the form (2) where G is uniquely determined by $\mathcal{L}$ up to a purely imaginary multiple of the identity operator.

Theorem 8 can be restated in the index free form ([24] Thm. 30.12).

Theorem 9 *Let $\mathcal{L}$ be the generator of a uniformly continuous QMS on $\mathcal{B}(\mathfrak{h})$, then there exist an Hilbert space $\mathfrak{k}$, a bounded linear operator $L : \mathfrak{h} \rightarrow \mathfrak{h} \otimes \mathfrak{k}$ and a bounded self-adjoint operator H in $\mathfrak{h}$ satisfying the following:*

1. $\mathcal{L}(x) = i[H, x] - \frac{1}{2} (L^* L x - 2L^*(x \otimes \mathbb{1}_{\mathfrak{k}})L + xL^*L)$ for all $x \in \mathcal{B}(\mathfrak{h})$;
2. the set $\{(x \otimes \mathbb{1}_{\mathfrak{k}})Lu : x \in \mathcal{B}(\mathfrak{h}), u \in \mathfrak{h}\}$ is total in $\mathfrak{h} \otimes \mathfrak{k}$.

Proof. Let $\mathfrak{k}$ be a Hilbert space with Hilbertian dimension equal to the length of the sequence $(L_k)_k$ and let (f_k) be an orthonormal basis of $\mathfrak{k}$. Defining $Lu = \sum_k L_k u \otimes f_k$, where (f_k) is an orthonormal basis of $\mathfrak{k}$ and the L_k are as in Theorem 8, a simple calculation shows that 1 is fulfilled.

Suppose that there exists a non-zero vector ξ orthogonal to the set of $(x \otimes \mathbb{1}_{\mathfrak{k}})Lu$ with $x \in \mathcal{B}(\mathfrak{h})$, $u \in \mathfrak{h}$; then $\xi = \sum_k v_k \otimes f_k$ with $v_k \in \mathfrak{h}$ and

$$0 = \langle \xi, (x \otimes \mathbb{1}_{\mathfrak{k}})Lu \rangle = \sum_k \langle v_k, xL_k u \rangle = \sum_k \langle L_k^* x^* v_k, u \rangle$$

for all $x \in \mathcal{B}(\mathfrak{h})$, $u \in \mathfrak{h}$. Hence, $\sum_k L_k^* x^* v_k = 0$. Since $\xi \neq 0$, we can suppose $\|v_1\| = 1$; then, putting $p = |v_1\rangle\langle v_1|$ and $x = py^*$, $y \in \mathcal{B}(\mathfrak{h})$, we get

$$0 = L_1^* y v_1 + \sum_{k \geq 2} \langle v_1, v_k \rangle L_k^* y v_1 = \left(L_1^* + \sum_{k \geq 2} \langle v_1, v_k \rangle L_k^* \right) y v_1. \quad (7)$$

Since $y \in \mathcal{B}(\mathfrak{h})$ is arbitrary, equation (7) contradicts the linear independence of the L_k 's. Therefore the set in (2) must be total. $\square$

The Hilbert space $\mathfrak{k}$ is called the *multiplicity space* of the completely positive part of $\mathcal{L}$. A unitary matrix $(u_{\ell j})_{\ell, j \geq 1}$, in the above basis $(f_k)_{k \geq 1}$,

clearly defines a unitary operator on $\mathbf{k}$. From now on we shall identify such matrices with operators on $\mathbf{k}$.

We end this section by establishing the relationship between the operators G, L_ℓ and G', L'_ℓ in two special GKSL representations of $\mathcal{L}$ and $\mathcal{L}'$ when these generator are both bounded.

The dual QMS $\mathcal{T}'$ clearly satisfies

$$\rho^{1/2} \mathcal{T}'_t(x) \rho^{1/2} = \mathcal{T}_{*t}(\rho^{1/2} x \rho^{1/2})$$

where $\mathcal{T}_*$ denotes the predual semigroup of $\mathcal{T}$. Since $\mathcal{L}'$ is bounded, differentiating at $t = 0$, we find the relationship among the generator $\mathcal{L}'$ of $\mathcal{T}$ and $\mathcal{L}_*$ of the predual semigroup $\mathcal{T}_*$ of $\mathcal{T}$

$$\rho^{1/2} \mathcal{L}'(x) \rho^{1/2} = \mathcal{L}_*(\rho^{1/2} x \rho^{1/2}). \quad (8)$$

Proposition 10 *Let $\mathcal{L}(a) = G^*a + aG + \sum_\ell L_\ell^* a L_\ell$ be a special GKSL representation of $\mathcal{L}$ with respect to a $\mathcal{T}$ -invariant state $\rho = \sum_k \rho_k |e_k\rangle\langle e_k|$. Then*

$$G^*u = \sum_{k \geq 1} \rho_k \mathcal{L}(|u\rangle\langle e_k|) e_k - \text{tr}(\rho G)u \quad (9)$$

$$Gv = \sum_{k \geq 1} \rho_k \mathcal{L}_*(|v\rangle\langle e_k|) e_k - \text{tr}(\rho G^*)v \quad (10)$$

for every $u, v \in \mathbf{h}$.

Proof. Since $\mathcal{L}(|u\rangle\langle v|) = |G^*u\rangle\langle v| + |u\rangle\langle Gv| + \sum_\ell |L_\ell^*u\rangle\langle L_\ell^*v|$, putting $v = e_k$ we have $G^*u = |G^*u\rangle\langle e_k| e_k$ and

$$G^*u = \mathcal{L}(|u\rangle\langle e_k|) e_k - \sum_\ell \langle e_k, L_\ell e_k \rangle L_\ell^* u - \langle e_k, G e_k \rangle u.$$

Multiplying both sides by ρ_k and summing on k , we find then

$$\begin{aligned} G^*u &= \sum_{k \geq 1} \rho_k \mathcal{L}(|u\rangle\langle e_k|) e_k - \sum_{\ell, k} \rho_k \langle e_k, L_\ell e_k \rangle L_\ell^* u - \sum_{k \geq 1} \rho_k \langle e_k, G e_k \rangle u \\ &= \sum_{k \geq 1} \rho_k \mathcal{L}(|u\rangle\langle e_k|) e_k - \sum_\ell \text{tr}(\rho L_\ell) L_\ell^* u - \text{tr}(\rho G)u \end{aligned}$$

and (9) follows since $\text{tr}(\rho L_j) = 0$. The identity (10) is now immediate computing the adjoint of G . $\square$

Proposition 11 *Let $\mathcal{T}'$ be the dual of a QMS $\mathcal{T}$ generated by $\mathcal{L}$ with normal invariant state ρ . If G and G' are the operators (10) in two GKSL representations of $\mathcal{L}$ and $\mathcal{L}'$ then*

$$G' \rho^{1/2} = \rho^{1/2} G^* + (\text{tr}(\rho G) - \text{tr}(\rho G')) \rho^{1/2}. \quad (11)$$

Moreover, we have $\text{tr}(\rho G) - \text{tr}(\rho G') = ic$ for some $c \in \mathbb{R}$.

Proof. The identities (10) and (8) yield

$$\begin{aligned}
G' \rho^{1/2} &= \sum_{k \geq 1} \mathcal{L}'_*(\rho^{1/2} |v\rangle \langle \rho_k^{1/2} e_k|) \rho_k^{1/2} e_k - \text{tr}(\rho G'^*) \rho^{1/2} v \\
&= \sum_{k \geq 1} \mathcal{L}'_*(\rho^{1/2} (|v\rangle \langle e_k|) \rho^{1/2}) \rho^{1/2} e_k - \text{tr}(\rho G'^*) \rho^{1/2} v \\
&= \sum_{k \geq 1} \rho^{1/2} \mathcal{L}(|v\rangle \langle e_k|) \rho^{1/2} \rho^{1/2} e_k - \text{tr}(\rho G'^*) \rho^{1/2} v \\
&= \rho^{1/2} G^* v + (\text{tr}(\rho G) - \text{tr}(\rho G'^*)) \rho^{1/2} v.
\end{aligned}$$

Therefore, we obtain (11). Right multiplying this equation by $\rho^{1/2}$ we have $G' \rho = \rho^{1/2} G^* \rho^{1/2} + (\text{tr}(\rho G) - \text{tr}(\rho G'^*)) \rho$, and, taking the trace,

$$\begin{aligned}
\text{tr}(\rho G) - \text{tr}(\rho G'^*) &= \text{tr}(G' \rho) - \text{tr}(\rho^{1/2} G^* \rho^{1/2}) \\
&= \text{tr}(G' \rho) - \text{tr}(G^* \rho) = -\overline{(\text{tr}(\rho G) - \text{tr}(\rho G'^*))};
\end{aligned}$$

this proves the last claim. $\square$

We can now prove as in [11] Th. 7.2 p. 358 the following

Theorem 12 *For all special GKSL representation of $\mathcal{L}$ by means of operators G, L_ℓ as in (1) there exists a special GKSL representation of $\mathcal{L}'$ by means of operators G', L'_ℓ such that:*

1. $G' \rho^{1/2} = \rho^{1/2} G^* + ic \rho^{1/2}$ for some $c \in \mathbb{R}$,
2. $L'_\ell \rho^{1/2} = \rho^{1/2} L_\ell^*$.

Proof. Since $\mathcal{L}'$ is bounded, it admits a special GKSL representation $\mathcal{L}'(a) = G'^* a + \sum_k L_k'^* a L'_k + a G'$. Moreover, by Proposition 11 we have $G' \rho^{1/2} = \rho^{1/2} G^* + ic$, $c \in \mathbb{R}$, and so (8) implies

$$\sum_k \rho^{1/2} L_k'^* x L'_k \rho^{1/2} = \sum_k L_k \rho^{1/2} x \rho^{1/2} L_k^*. \quad (12)$$

Let $\mathbf{k}$ (resp. $\mathbf{k}'$) be the multiplicity space of the completely positive part of $\mathcal{L}$ (resp. $\mathcal{L}'$) and define an operator $X : \mathbf{h} \otimes \mathbf{k}' \rightarrow \mathbf{h} \otimes \mathbf{k}$

$$X(x \otimes \mathbb{1}_{\mathbf{k}'}) L' \rho^{1/2} u = (x \otimes \mathbb{1}_{\mathbf{k}}) (\rho^{1/2} \otimes \mathbb{1}_{\mathbf{k}}) L^* u$$

for all $x \in \mathcal{B}(\mathbf{h})$ and $u \in \mathbf{h}$, where $L : \mathbf{h} \rightarrow \mathbf{h} \otimes \mathbf{k}$, $Lu = \sum_k L_k u \otimes f_k$, $L' : \mathbf{h} \rightarrow \mathbf{h} \otimes \mathbf{k}'$, $L'u = \sum_k L'_k u \otimes f'_k$, $(f_k)_k$ and $(f'_k)_k$ are orthonormal bases of $\mathbf{k}$ and $\mathbf{k}'$ respectively. Thus, by (12),

$$\begin{aligned}
\langle X(x \otimes \mathbb{1}_{\mathbf{k}'}) L' \rho^{1/2} u, X(y \otimes \mathbb{1}_{\mathbf{k}'}) L' \rho^{1/2} v \rangle &= \sum_k \langle u, \rho^{1/2} L_k'^* x^* y L'_k \rho^{1/2} v \rangle \\
&= \langle (x \otimes \mathbb{1}_{\mathbf{k}'}) L' \rho^{1/2} u, (y \otimes \mathbb{1}_{\mathbf{k}'}) L' \rho^{1/2} v \rangle
\end{aligned}$$

for all $x, y \in \mathcal{B}(\mathfrak{h})$ and $u, v \in \mathfrak{h}$, i.e. X preserves the scalar product. Therefore, since the set $\{(x \otimes \mathbb{1}_{\mathfrak{k}'})L'\rho^{1/2}u \mid x \in \mathcal{B}(\mathfrak{h}), u \in \mathfrak{h}\}$ is total in $\mathfrak{h} \otimes \mathfrak{k}'$ (for $\rho^{1/2}(\mathfrak{h})$ is dense in $\mathfrak{h}$ and Theorem 9 holds), we can extend X to an unitary operator from $\mathfrak{h} \otimes \mathfrak{k}'$ to $\mathfrak{h} \otimes \mathfrak{k}$. As a consequence we have $X^*X = \mathbb{1}_{\mathfrak{h} \otimes \mathfrak{k}'}$.

Moreover, since $X(y \otimes \mathbb{1}_{\mathfrak{k}'}) = (y \otimes \mathbb{1}_{\mathfrak{k}})X$ for all $y \in \mathcal{B}(\mathfrak{h})$, we can conclude that $X = \mathbb{1}_{\mathfrak{h}} \otimes Y$ for some unitary map $Y : \mathfrak{k}' \rightarrow \mathfrak{k}$.

The definition of X implies then

$$(\rho^{1/2} \otimes \mathbb{1}_{\mathfrak{k}})L^* = XL'\rho^{1/2} = (\mathbb{1}_{\mathfrak{h}} \otimes Y)L'\rho^{1/2}.$$

This means that, replacing L' by $(\mathbb{1}_{\mathfrak{h}} \otimes Y)L'$, or more precisely L'_k by $\sum_{\ell} u_{k\ell}L'_\ell$ for all k , we have

$$\rho^{1/2}L_k^* = L'_k\rho^{1/2}.$$

Since $\text{tr}(\rho L'_k) = \text{tr}(\rho L_k^*) = 0$ and, from $\mathcal{L}'(\mathbb{1}) = 0$, $G'^* + G' = -\sum_k L_k'^* L'_k$, the properties of a special GKSL representation follow. $\square$

Remark 3 Condition 2 implies that the completely positive parts $\Phi(x) = \sum_{\ell} L_{\ell}^* x L_{\ell}$ and Φ' of the generators $\mathcal{L}$ and $\mathcal{L}'$, respectively are mutually adjoint i.e.

$$\text{tr}(\rho^{1/2}\Phi'(x)\rho^{1/2}y) = \text{tr}(\rho^{1/2}x\rho^{1/2}\Phi(y)) \quad (13)$$

for all $x, y \in \mathcal{B}(\mathfrak{h})$. As a consequence, also the maps $x \rightarrow G^*x + xG$ and $x \rightarrow (G')^*x + xG'$ are mutually adjoint.

4 Generators of Standard Detailed Balance QMSs

In this section we characterise the generators of norm-continuous QMSs satisfying the SQDB of Definition 4.

We start noting that, since ρ is invariant for $\mathcal{T}$ and $\mathcal{T}'$, i.e. $\mathcal{L}_*(\rho) = \mathcal{L}'_*(\rho) = 0$, the operator K commutes with ρ . Moreover, by comparing two special GKSL representations of $\mathcal{L}$ and $\mathcal{L}' + 2i[K, \cdot]$, we have immediately the following

Lemma 13 *A QMS $\mathcal{T}$ satisfies the SQDB $\mathcal{L} - \mathcal{L}' = 2i[K, \cdot]$ if and only if for all special GKSL representations of the generators $\mathcal{L}$ and $\mathcal{L}'$ by means of operators G, L_k and G', L'_k respectively, we have*

$$G = G' + 2iK + ic \quad L'_k = \sum_j u_{kj}L_j$$

for some $c \in \mathbb{R}$ and some unitary $(u_{kj})_{kj}$ on $\mathfrak{k}$.

Since we know the relationship between the operators G', L'_k and G, L_k thanks to Theorem 12, we can now characterise generators of QMSs satisfying the SQDB. We emphasize the following definition of T -symmetric matrix (operator) on $\mathbf{k}$ in order to avoid confusion with the usual notion of symmetric operator X meaning that X^* is an extension of X .

Definition 14 Let $Y = (y_{k\ell})_{k,\ell \geq 1}$ be a matrix with entries indexed by k, ℓ running on the set (finite or infinite) of indices of the sequence $(L_\ell)_{\ell \geq 1}$. We denote by Y^T the transpose matrix $Y^T = (y_{\ell k})_{k,\ell \geq 1}$. The matrix Y is called T -symmetric if $Y = Y^T$.

Theorem 15 $\mathcal{T}$ satisfies the SQDB if and only if for all special GKSL representation of the generator $\mathcal{L}$ by means of operators G, L_k there exists a T -symmetric unitary $(u_{m\ell})_{m\ell}$ on $\mathbf{k}$ such that

$$\rho^{1/2} L_k^* = \sum_{\ell} u_{k\ell} L_{\ell} \rho^{1/2}, \quad (14)$$

for all $k \geq 1$.

Proof. Given a special GKSL of $\mathcal{L}$, Theorem 12 allows us to write the dual $\mathcal{L}'$ in a special GKSL representation by means of operators G', L'_k with

$$G' \rho^{1/2} = \rho^{1/2} G^*, \quad L'_k \rho^{1/2} = \rho^{1/2} L_k^*. \quad (15)$$

Suppose first that $\mathcal{T}$ satisfies the SQDB. Since $L'_k = \sum_j u_{kj} L_j$ for some unitary $(u_{kj})_{kj}$ by Lemma 13, we can find (14) substituting L'_k with $\sum_j u_{kj} L_j$ in the second formula (15).

Finally we show that the unitary matrix $u = (u_{m\ell})_{m\ell}$ is T -symmetric. Indeed, taking the adjoint of (14) we find $L_{\ell} \rho^{1/2} = \sum_m \bar{u}_{\ell m} \rho^{1/2} L_m^*$. Writing $\rho^{1/2} L_m^*$ as in (14) we have then

$$L_{\ell} \rho^{1/2} = \sum_{m,k} \bar{u}_{\ell m} u_{mk} L_k \rho^{1/2} = \sum_k ((u^*)^T u)_{\ell k} L_k \rho^{1/2}.$$

The operators $L_{\ell} \rho^{1/2}$ are linearly independent by property (iii) Theorem 8 of a special GKSL representation, therefore $(u^*)^T u$ is the identity operator on $\mathbf{k}$. Since u is also unitary, we have also $u^* u = (u^*)^T u$, namely $u^* = (u^*)^T$ and $u = u^T$.

Conversely, if (14) holds, by (15), we have $L'_k \rho^{1/2} = \sum_{\ell} u_{k\ell} L_{\ell} \rho^{1/2}$, so that $L'_k = \sum_{\ell} u_{k\ell} L_{\ell}$ for all k and for some unitary $(u_{kj})_{kj}$. Therefore, thanks to Lemma 13, to conclude it is enough to prove that $G = G' + i(2K + c)$ namely, that $G - G'$ is anti self-adjoint.

To this end note that, since ρ is an invariant state, we have

$$0 = \rho G^* + \sum_k L_k \rho L_k^* + G \rho, \quad (16)$$

with

$$\begin{aligned} \sum_k L_k \rho L_k^* &= \sum_k (L_k \rho^{1/2})(\rho^{1/2} L_k^*) = \sum_k \sum_{\ell, j} \bar{u}_{k\ell} u_{kj} \rho^{1/2} L_\ell^* L_j \rho^{1/2} \\ &= \sum_\ell \rho^{1/2} L_\ell^* L_\ell \rho^{1/2} = -\rho^{1/2} (G + G^*) \rho^{1/2}, \end{aligned}$$

(for condition (14) holds) and so, by substituting in equation (16) we get

$$\begin{aligned} 0 &= \rho G^* - \rho^{1/2} G \rho^{1/2} - \rho^{1/2} G^* \rho^{1/2} + G \rho = \rho^{1/2} (\rho^{1/2} G^* - G \rho^{1/2}) \\ &\quad - (\rho^{1/2} G^* - G \rho^{1/2}) \rho^{1/2} = [G \rho^{1/2} - \rho^{1/2} G^*, \rho^{1/2}], \end{aligned}$$

i.e. $G \rho^{1/2} - \rho^{1/2} G^*$ commutes with $\rho^{1/2}$.

We can now prove that $G - G'$ is anti self-adjoint. Clearly, it suffices to show that $\rho^{1/2} G \rho^{1/2} - \rho^{1/2} G' \rho^{1/2}$ is anti self-adjoint. Indeed, by (15), we have

$$\begin{aligned} (\rho^{1/2} G \rho^{1/2} - \rho^{1/2} G' \rho^{1/2})^* &= (\rho^{1/2} G \rho^{1/2} - \rho G^*)^* \\ &= (\rho^{1/2} (G \rho^{1/2} - \rho^{1/2} G^*))^* \\ &= ((G \rho^{1/2} - \rho^{1/2} G^*) \rho^{1/2})^* \\ &= \rho G^* - \rho^{1/2} G \rho^{1/2} = \rho^{1/2} G' \rho^{1/2} - \rho^{1/2} G \rho^{1/2} \end{aligned}$$

because $G \rho^{1/2} - \rho^{1/2} G^*$ commutes with $\rho^{1/2}$. This completes the proof. $\square$

It is worth noticing that, as in Remark 3, $\mathcal{T}$ satisfies the SQDB if and only if the completely positive part Φ of the generator $\mathcal{L}$ is symmetric. This improves our previous result, Thm. 7.3 [11], where we gave $G \rho^{1/2} = \rho^{1/2} G^* - (2iK + ic) \rho^{1/2}$ for some $c \in \mathbb{R}$ as an additional condition. Here we showed that it follows from (14) and the invariance of ρ .

Remark 4 Note that (14) holds for the operators L_ℓ of a special GKSL representation of $\mathcal{L}$ if and only if it is true for *all* special GKSL representations because of the second part of Theorem 8. Therefore the conclusion of Theorem 15 holds true also if and only if we can find a single special GKSL representation of $\mathcal{L}$ satisfying (14).

The T -symmetric unitary $(u_{m\ell})_{m\ell}$ is determined by the L_ℓ 's because they are linearly independent. We shall now exploit this fact to give a more geometrical characterisation of SQDB.

When the SQDB holds, the matrices $(b_{kj})_{k,j \geq 1}$ and $(c_{kj})_{k,j \geq 1}$ with

$$b_{kj} = \text{tr}(\rho^{1/2} L_k^* \rho^{1/2} L_j^*), \quad \text{and} \quad c_{kj} = \text{tr}(\rho L_k^* L_j) \quad (17)$$

define two trace class operators B and C on $\mathbf{k}$ by Lemma 26 (see the Appendix); B is T -symmetric and C is self-adjoint. Moreover, it admits a self-adjoint inverse C^{-1} because ρ is faithful. When $\mathbf{k}$ is infinite dimensional, C^{-1} is unbounded and its domain coincides with the range of C .

We can now give the following characterisation of QMS satisfying the SQDB condition which is more direct because the unitary $(u_{k\ell})_{k\ell}$ in Theorem 15 is explicitly given by $C^{-1}B$.

Theorem 16 *$\mathcal{T}$ satisfies the SQDB if and only if the operators G, L_k of a special GKSL representation of the generator $\mathcal{L}$ satisfy the following conditions:*

- (i) *the closed linear span of $\{\rho^{1/2} L_\ell^* \mid \ell \geq 1\}$ and $\{L_\ell \rho^{1/2} \mid \ell \geq 1\}$ in the Hilbert space of Hilbert-Schmidt operators on $\mathbf{h}$ coincide,*
- (ii) *the trace-class operators B, C defined by (17) satisfy $CB = BC^T$ and $C^{-1}B$ is unitary T -symmetric.*

Proof. If $\mathcal{T}$ satisfies the SQDB then, by Theorem 15, the identity (14) holds. The series in the right-hand side of (14) is convergent with respect to the Hilbert-Schmidt norm because

$$\begin{aligned} \left\| \sum_{m+1 \leq \ell \leq n} u_{k\ell} L_\ell \rho^{1/2} \right\|_{HS}^2 &= \sum_{m+1 \leq \ell, \ell' \leq n} \bar{u}_{k\ell'} u_{k\ell} \text{tr}(\rho L_{\ell'}^* L_\ell) \\ &\leq \frac{1}{2} \sum_{m+1 \leq \ell, \ell' \leq n} |u_{k\ell'}|^2 |u_{k\ell}|^2 + \frac{1}{2} \sum_{m+1 \leq \ell, \ell' \leq n} |c_{\ell'\ell}|^2 \\ &\leq \frac{1}{2} \left(\sum_{m+1 \leq \ell \leq n} |u_{k\ell}|^2 \right)^2 + \frac{1}{2} \sum_{m+1 \leq \ell, \ell' \leq n} |c_{\ell'\ell}|^2 \end{aligned}$$

and the right-hand side vanishes as n, m go to infinity because the operator C is trace-class by Lemma 26 and the columns of $U = (u_{k\ell})_{k\ell}$ are unit vectors in $\mathbf{k}$ by unitarity.

Left multiplying both sides of (14) by $\rho^{1/2} L_j^*$ and taking the trace we find $B = CU^T = CU$. It follows that the range of the operators B, CU and C coincide and $C^{-1}B = U$ is everywhere defined, unitary and T -symmetric

because U is T -symmetric. Moreover, since B is T -symmetric by the cyclic property of the trace, we have also

$$BC^T = CU^T C^T = C(CU)^T = CB^T = CB.$$

Conversely, we show that (i) and (ii) imply the SQDB. To this end notice that, by the spectral theorem we can find a unitary linear transformation $V = (v_{mn})_{m,n \geq 1}$ on $\mathfrak{h}$ such that V^*CV is diagonal. Therefore, choosing a new GKSL of the generator $\mathcal{L}$ by means of the operators $L''_k = \sum_{n \geq 1} v_{nk} L_n$, if necessary, we can suppose that both $(L_\ell \rho^{1/2})_{\ell \geq 1}$ and $(\rho^{1/2} L_k^*)_{k \geq 1}$ are *orthogonal* bases of the same closed linear space. Note that

$$\mathrm{tr}(\rho^{1/2} (L'')^*_k \rho^{1/2} (L'')^*_j) = \sum_{m,n \geq 1} \bar{v}_{nk} \bar{v}_{mj} \mathrm{tr}(\rho^{1/2} L_n^* \rho^{1/2} L_m^*)$$

and the operator B , after this change of GKSL representation, becomes $V^*B(V^*)^T$ which is also T -symmetric.

Writing the expansion of $\rho^{1/2} L_k^*$ with respect to the orthogonal basis $(L_\ell \rho^{1/2})_{\ell \geq 1}$, for all $k \geq 1$ we have

$$\rho^{1/2} L_k^* = \sum_{\ell \geq 1} \frac{\mathrm{tr}(\rho^{1/2} L_\ell^* \rho^{1/2} L_k^*)}{\|L_\ell \rho^{1/2}\|_{HS}^2} L_\ell \rho^{1/2}. \quad (18)$$

In this way we find a matrix Y of complex numbers $y_{k\ell}$ such that $\rho^{1/2} L_k^* = \sum_{\ell} y_{k\ell} L_\ell \rho^{1/2}$ and the series is Hilbert-Schmidt norm convergent. Clearly, since C is diagonal and B is T -symmetric, $y_{k\ell} = (BC^{-1})_{k\ell} = ((B(C^{-1})^T)_{k\ell} = ((C^{-1}B)^T)_{k\ell}$. It follows from (ii) that Y coincides with the unitary operator $(C^{-1}B)^T$ and (14) holds. Moreover, Y is symmetric because

$$y_{\ell k} = (BC^{-1})_{\ell k} = ((B(C^{-1})^T)_{\ell k} = (C^{-1}B)_{k\ell} = y_{k\ell}.$$

This completes the proof. $\square$

Formula (18) has the following consequence.

Corollary 17 *Suppose that a QMS $\mathcal{T}$ satisfies the SQDB condition. For every special GKSL representation of $\mathcal{L}$ with operators $L_\ell \rho^{1/2}$ that are orthogonal in the Hilbert space of Hilbert-Schmidt operators on $\mathfrak{h}$ if $\mathrm{tr}(\rho^{1/2} L_\ell^* \rho^{1/2} L_k^*) \neq 0$ for a pair of indices $k, \ell \geq 1$, then $\mathrm{tr}(\rho L_\ell^* L_\ell) = \mathrm{tr}(\rho L_k^* L_k)$.*

Proof. It suffices to note that the matrix $(u_{k\ell})$ with entries

$$u_{k\ell} = \frac{\mathrm{tr}(\rho^{1/2} L_\ell^* \rho^{1/2} L_k^*)}{\|L_\ell \rho^{1/2}\|_{HS}^2} = \frac{\mathrm{tr}(\rho^{1/2} L_\ell^* \rho^{1/2} L_k^*)}{\mathrm{tr}(\rho L_\ell^* L_\ell)}$$

must be T -symmetric. $\square$

Remark 5 The matrix C can be viewed as the covariance matrix of the zero-mean (recall that $\text{tr}(\rho L_\ell) = 0$) “random variables” $\{L_\ell \mid \ell \geq 1\}$ and in a similar way, B can be viewed as a sort of mixed covariance matrix between the previous random variable and the adjoint $\{L_\ell^* \mid \ell \geq 1\}$. Thus the SQDB condition holds when the random variables L_ℓ right multiplied by $\rho^{1/2}$ and the adjoint variables L_ℓ^* left multiplied by $\rho^{1/2}$ generate the same subspace of Hilbert-Schmidt operators and the mixed covariance matrix B is a left unitary transformation of the covariance matrix C .

If we consider a special GKSL representation of $\mathcal{L}$ with operators $L_\ell \rho^{1/2}$ that are orthogonal, then, by Corollary 17 and the identity $\|L_\ell \rho^{1/2}\|_{HS} = \|L_k \rho^{1/2}\|_{HS}$, the unitary matrix U can be written as $C^{-1/2} B C^{-1/2}$. This, although not positive definite, can be interpreted as a correlation coefficient matrix of $\{L_\ell \mid \ell \geq 1\}$ and $\{L_\ell^* \mid \ell \geq 1\}$.

The characterisation of generators of symmetric QMSs with respect to the $s = 1/2$ scalar product follows along the same lines.

Theorem 18 *A norm-continuous QMS $\mathcal{T}$ is symmetric if and only if there exists a special GKSL representation of the generator $\mathcal{L}$ by means of operators G, L_ℓ such that*

- (1) $G \rho^{1/2} = \rho^{1/2} G^* + i c \rho^{1/2}$ for some $c \in \mathbb{R}$,
- (2) $\rho^{1/2} L_k^* = \sum_\ell u_{k\ell} L_\ell \rho^{1/2}$, for all k , for some unitary $(u_{k\ell})_{k\ell}$ on $\mathbf{k}$ which is also T -symmetric.

Proof. Choose a special GKSL representation of $\mathcal{L}$ by means of operators G, L_k . Theorem 12 allows us to write the symmetric dual $\mathcal{L}'$ in a special GKSL representation by means of operators G', L'_k as in (15).

Suppose first that $\mathcal{T}$ is KMS-symmetric. Comparing the special GKSL representations of $\mathcal{L}$ and $\mathcal{L}'$, by Theorem 8 we find

$$G = G' + i c, \quad L'_k = \sum_j u_{kj} L_j,$$

for some unitary matrix (u_{kj}) and some $c \in \mathbb{R}$. This, together with (15) implies that conditions (1) and (2) hold.

Assume now that conditions (1) and (2) hold. Taking the adjoint of (2) we find immediately $L_k \rho^{1/2} = \sum_k \bar{u}_{k\ell} \rho^{1/2} L_\ell^*$. Then straightforward computation,

by the unitarity of the matrix $(u_{k\ell})$, yields

$$\begin{aligned}
\mathcal{L}_*(\rho^{1/2}x\rho^{1/2}) &= G\rho^{1/2}x\rho^{1/2} + \sum_k L_k\rho^{1/2}x\rho^{1/2}L_k^* + \rho^{1/2}x\rho^{1/2}G^* \\
&= \rho^{1/2}G^*x\rho^{1/2} + \sum_{\ell,k,j} \bar{u}_{k\ell} u_{kj} \rho^{1/2}L_k^*xL_j\rho^{1/2} + \rho^{1/2}xG\rho^{1/2} \\
&= \rho^{1/2}\mathcal{L}(x)\rho^{1/2}
\end{aligned}$$

for all $x \in \mathcal{B}(\mathfrak{h})$. Iterating we find $\mathcal{L}_*(\rho^{1/2}x\rho^{1/2}) = \rho^{1/2}\mathcal{L}^n(x)\rho^{1/2}$ for all $n \geq 0$, therefore, exponentiating, we find $\mathcal{T}_{*t}(\rho^{1/2}x\rho^{1/2}) = \rho^{1/2}\mathcal{T}_t(x)\rho^{1/2}$ for all $t \geq 0$. This, together with (3), implies that $\mathcal{T}$ is KMS-symmetric. $\square$

Remark 6 Note that condition (2) in Theorem 18 implies that the completely positive part of $\mathcal{L}$ is KMS-symmetric. This makes a parallel with Theorem 12 where condition (2) implies that the completely positive parts of the generators $\mathcal{L}$ and $\mathcal{L}'$ are mutually adjoint.

The above theorem simplifies a previous result by Park ([23] Thm 2.2) where conditions (1) and (2) appear in a much more complicated way.

5 Generators of Standard Detailed Balance (with time reversal) QMSs

We shall now study generators of semigroups satisfying the SQDB- θ introduced in Definition 5 involving the time reversal operation. In this section, we always assume that the invariant state ρ and the anti-unitary time reversal θ commute.

The relationship between the QMS satisfying the SQDB- θ , its dual and their generators is clarified by the following

Proposition 19 *A QMS $\mathcal{T}$ satisfies the SQDB- θ if and only if the dual semigroup $\mathcal{T}'$ is given by*

$$\mathcal{T}'_t(x) = \theta\mathcal{T}_t(\theta x\theta)\theta \quad \text{for all } x \in \mathcal{B}(\mathfrak{h}). \quad (19)$$

In particular, if $\mathcal{T}$ is norm-continuous, then $\mathcal{T}'$ is also norm-continuous. Moreover, in this case $\mathcal{T}'$ is generated by

$$\mathcal{L}'(x) = \theta\mathcal{L}(\theta x\theta)\theta, \quad x \in \mathcal{B}(\mathfrak{h}). \quad (20)$$

Proof. Suppose that $\mathcal{T}$ satisfies the SQDB- θ and put $\sigma(x) = \theta x \theta$. Taking $t = 0$ equation (6) reduces to $\text{tr}(\rho^{1/2} x \rho^{1/2} y) = \text{tr}(\rho^{1/2} \sigma(y^*) \rho^{1/2} \sigma(x^*))$ for all $x, y \in \mathcal{B}(\mathfrak{h})$, so that

$$\begin{aligned} \text{tr}(\rho^{1/2} x \rho^{1/2} \mathcal{T}_t(y)) &= \text{tr}(\rho^{1/2} \sigma(y^*) \rho^{1/2} \mathcal{T}_t(\sigma(x^*))) \\ &= \text{tr}(\rho^{1/2} \sigma(\mathcal{T}_t(\sigma(x^*))^* \rho^{1/2} \sigma(\sigma(y^*)^*)) \\ &= \text{tr}(\rho^{1/2} \sigma(\mathcal{T}_t(\sigma(x))) \rho^{1/2} y) \end{aligned}$$

for every $x, y \in \mathcal{B}(\mathfrak{h})$ and (19) follows. Therefore, if $\mathcal{T}$ is norm continuous, $\mathcal{T}'_t = (\sigma \circ \mathcal{T}_t \circ \sigma)_t$ is also.

Conversely, if (19) holds, the commutation between ρ and θ implies

$$\begin{aligned} \text{tr}(\rho^{1/2} \mathcal{T}'_t(x) \rho^{1/2} y) &= \text{tr}(\rho^{1/2} \theta \mathcal{T}_t(\theta x \theta) \theta \rho^{1/2} y) \\ &= \text{tr}(\theta (\rho^{1/2} \mathcal{T}_t(\theta x \theta) \theta \rho^{1/2} y \theta) \theta) \\ &= \text{tr}(\rho^{1/2} \theta y^* \rho^{1/2} \theta \mathcal{T}_t(\theta x^* \theta)) \end{aligned}$$

and (19) is proved. Now (20) follows from (19) differentiating at $t = 0$. $\square$

We can now describe the relationship between special GKSL representations of $\mathcal{L}$ and $\mathcal{L}'$.

Proposition 20 *If $\mathcal{T}$ satisfies the SQDB- θ then, for every special GKSL representation of $\mathcal{L}$ by means of operators H, L_k , the operators $H' = -\theta H \theta$ and $L'_k = \theta L_k \theta$ yield a special GKSL representation of $\mathcal{L}'$.*

Proof. Consider a special GKSL representation of $\mathcal{L}$ by means of operators H, L_k . Since $\mathcal{L}'(a) = \theta \mathcal{L}(\theta a \theta) \theta$ by Proposition 19, from the antilinearity of θ and $\theta^2 = \mathbb{1}$ we get

$$\begin{aligned} \theta \mathcal{L}'(a) \theta &= i[H, \theta a \theta] - \frac{1}{2} \sum_k (L_k^* L_k \theta a \theta - 2L_k^* \theta a \theta L_k + \theta a \theta L_k^* L_k) \\ &= i\theta (\theta H \theta a - a \theta H \theta) \theta + \sum_k \theta ((\theta L_k^* \theta) a (\theta L_k \theta)) \theta \\ &\quad - \frac{1}{2} \sum_k \theta ((\theta L_k^* \theta) (\theta L_k \theta) a + a (\theta L_k^* \theta) (\theta L_k \theta)) \theta \\ &= \theta (-i[\theta H \theta, a]) \theta - \frac{1}{2} \sum_k \theta (L_k'^* L_k' a - 2L_k'^* a L_k' + a L_k'^* L_k') \theta, \end{aligned}$$

where $L'_k := \theta L_k \theta$. Therefore, putting $H' = -\theta H \theta$, we find a GKSL representation of $\mathcal{L}'$ which is also special because $\text{tr}(\rho L'_k) = \text{tr}(\theta \rho L_k \theta) = \text{tr}(L_k^* \rho) = \overline{\text{tr}(\rho L_k)} = 0$. $\square$

The structure of generators of QMSs satisfying the SQDB- θ is described by the following

Theorem 21 *A QMS $\mathcal{T}$ satisfies the SQDB- θ condition if and only if there exists a special GKSL representation of $\mathcal{L}$, with operators G, L_ℓ , such that:*

1. $\rho^{1/2}\theta G^*\theta = G\rho^{1/2}$,
2. $\rho^{1/2}\theta L_k^*\theta = \sum_j u_{kj}L_j\rho^{1/2}$ for a self-adjoint unitary $(u_{kj})_{kj}$ on $\mathbf{k}$.

Proof. Suppose that $\mathcal{T}$ satisfies the SQDB- θ condition and consider a special GKSL representation of the generator $\mathcal{L}$ with operators G, L_k . The operators $-\theta H\theta$ and $\theta L_k\theta$ give then a special GKSL representation of $\mathcal{L}'$ by Proposition 20. Moreover, by Theorem 12, we have another special GKSL representation of $\mathcal{L}'$ by means of operators G', L'_k such that $G'\rho^{1/2} = \rho^{1/2}G^* + ic\rho^{1/2}$ for some $c \in \mathbb{R}$, and $L'_k\rho^{1/2} = \rho^{1/2}L_k^*$. Therefore there exists a unitary $(v_{kj})_{kj}$ on $\mathbf{k}$ such that $L'_k = \sum_j v_{kj}\theta L_j\theta$, and $\rho^{1/2}L_k^* = \sum_j v_{kj}\theta L_j\theta\rho^{1/2}$. Condition 2 follows then with $u_{kj} = \bar{v}_{kj}$ left and right multiplying by the antiunitary θ .

In order to find condition 1, first notice that by the unitarity of $(v_{kj})_{kj}$

$$\sum_k L_k'^* L'_k = \sum_k \theta L_k^* L_k \theta. \quad (21)$$

Now, by the uniqueness of G' up to a purely imaginary multiple of the identity in a special GKSL representation, $H' = (G'^* - G')/(2i)$ is equal to $-\theta H\theta + c_1$ for some $c_1 \in \mathbb{R}$. From (21) and $G'\rho^{1/2} = \rho^{1/2}G^* + ic\rho^{1/2}$ we obtain then

$$\begin{aligned} \rho^{1/2}G^* + ic\rho^{1/2} &= G'\rho^{1/2} = -iH'\rho^{1/2} - \frac{1}{2} \sum_k L_k'^* L'_k \rho^{1/2} \\ &= i\theta H\theta \rho^{1/2} + ic_1 \rho^{1/2} - \frac{1}{2} \sum_k \theta L_k^* L_k \theta \rho^{1/2} \\ &= \theta G\theta \rho^{1/2} + ic_1 \rho^{1/2}. \end{aligned}$$

It follows that $\rho^{1/2}\theta G^*\theta = G\rho^{1/2} + ic_2\rho^{1/2}$ for some $c_2 \in \mathbb{R}$. Left multiplying by $\rho^{1/2}$ and tracing we find

$$c_2 = \text{tr}(\theta\rho G^*\theta) - \text{tr}(\rho G) = \text{tr}(G\rho) - \text{tr}(\rho G) = 0$$

and condition 1 holds.

Finally we show that the square of the unitary $(u_{kj})_{kj}$ on $\mathbf{k}$ is the identity operator. Indeed, taking the adjoint of the identity $\rho^{1/2}\theta L_k^*\theta = \sum_j u_{kj}L_j\rho^{1/2}$, we have

$$\theta L_k\theta\rho^{1/2} = \sum_j \bar{u}_{kj}\rho^{1/2}L_j^*.$$

Left and right multiplying by the antilinear time reversal θ (commuting with ρ) we find

$$L_k \rho^{1/2} = \sum_j \theta \bar{u}_{kj} \rho^{1/2} L_j^* \theta = \sum_j u_{kj} \rho^{1/2} \theta L_j^* \theta$$

Writing $\rho^{1/2} \theta L_j^* \theta$ as $\sum_m u_{jm} L_m \rho^{1/2}$ by condition 2 we have then

$$L_k \rho^{1/2} = \sum_{j,m} u_{kj} u_{jm} L_m \rho^{1/2} = \sum_m (u^2)_{km} L_m \rho^{1/2}$$

which implies that $u^2 = \mathbb{1}$ by the linear independence of the $L_m \rho^{1/2}$. Therefore, since u is unitary, $u = u^*$.

Conversely, if 1 and 2 hold, we can write $\rho^{1/2} \theta \mathcal{L}(\theta x \theta) \theta \rho^{1/2}$ as

$$\begin{aligned} & \rho^{1/2} \theta G^* \theta x \rho^{1/2} + \sum_k \rho^{1/2} \theta L_k^* \theta x \theta L_k \theta \rho^{1/2} + \rho^{1/2} x \theta G \theta \rho^{1/2} \\ &= G \rho^{1/2} x \rho^{1/2} + \sum_j L_j \rho^{1/2} x \rho^{1/2} L_j^* + \rho^{1/2} x \rho^{1/2} G^*. \end{aligned}$$

This, by Theorem 12, can be written as

$$\rho^{1/2} (G')^* x \rho^{1/2} + \sum_j \rho^{1/2} (L_j')^* x L_j' \rho^{1/2} + \rho^{1/2} x G' \rho^{1/2} = \rho^{1/2} \mathcal{L}'(x) \rho^{1/2}$$

It follows that $\theta \mathcal{L}(\theta x \theta) \theta = \mathcal{L}'(x)$ for all $x \in \mathcal{B}(\mathfrak{h})$ because ρ is faithful. Moreover, it is easy to check by induction that $\theta \mathcal{L}^n(\theta x \theta) \theta = (\mathcal{L}')^n(x)$ for all $n \geq 0$. Therefore $\theta \mathcal{T}_t(\theta x \theta) \theta = \mathcal{T}'_t(x)$ for all $t \geq 0$ and $\mathcal{T}$ satisfies the SQDB- θ condition by Proposition 19. $\square$

We now provide a geometrical characterisation of the SQDB- θ condition as in Theorem 16. To this end we introduce the trace class operator R on $\mathfrak{k}$

$$R_{jk} = \text{tr} \left(\rho^{1/2} L_j^* \rho^{1/2} \theta L_k^* \theta \right) \quad (22)$$

A direct application of Lemma 26 shows that R is trace class. Moreover it is self-adjoint because, by the property $\text{tr}(\theta x \theta) = \text{tr}(x^*)$ of the antilinear time reversal, we have

$$\begin{aligned} \overline{R_{jk}} &= \overline{\text{tr} \left(\rho^{1/2} L_j^* \rho^{1/2} \theta L_k^* \theta \right)} \\ &= \text{tr} \left(\theta (L_k \theta \rho^{1/2} L_j \rho^{1/2} \theta) \theta \right) \\ &= \text{tr} \left(\rho^{1/2} \theta L_j^* \rho^{1/2} \theta L_k^* \right) \\ &= \text{tr} \left((\rho^{1/2} \theta L_j^* \theta) (\rho^{1/2} L_k^*) \right) = R_{kj}. \end{aligned}$$

Theorem 22 $\mathcal{T}$ satisfies the SQDB- θ if and only if the operators G, L_k of a special GKSL of the generator $\mathcal{L}$ fulfill the following conditions:

1. $\rho^{1/2}\theta G^*\theta = G\rho^{1/2}$,
2. the closed linear span of $\{\rho^{1/2}\theta L_\ell^*\theta \mid \ell \geq 1\}$ and $\{L_\ell\rho^{1/2} \mid \ell \geq 1\}$ in the Hilbert space of Hilbert-Schmidt operators on $\mathfrak{h}$ coincide,
3. the self-adjoint trace class operators R, C defined by (17) and (22) commute and $C^{-1}R$ is unitary and self-adjoint.

Proof. It suffices to show that conditions 2 and 3 above are equivalent to condition 2 of Theorem 21.

If $\mathcal{T}$ satisfies the SQBD- θ , then it can be shown as in the proof of Theorem 16 that 2 follows from condition 2 of Theorem 21. Moreover, left multiplying by $\rho^{1/2}L_\ell^*$ the identity $\rho^{1/2}\theta L_k^*\theta = \sum_j u_{kj}L_j\rho^{1/2}$ and tracing, we find

$$\mathrm{tr}(\rho^{1/2}L_\ell^*\rho^{1/2}\theta L_k^*\theta) = \sum_j u_{kj}\mathrm{tr}(\rho L_\ell^*L_j)$$

for all k, ℓ i.e. $R = CU^T$. The operator U^T is also self-adjoint and unitary. Therefore R and C have the same range and, since the domain of C^{-1} coincides with the range of C , the operator $C^{-1}R$ is everywhere defined, unitary and self-adjoint. It follows that the densely defined operator RC^{-1} is a restriction of $(C^{-1}R)^* = C^{-1}R$ and $CR = RC$.

In order to prove, conversely, that 2 and 3 imply condition 2 of Theorem 21, we first notice that, by the spectral theorem there exists a unitary $V = (v_{mn})_{m,n \geq 1}$ on the multiplicity space $\mathfrak{k}$ such that V^*CV is diagonal. Choosing a new GKSL representation of the generator $\mathcal{L}$ by means of the operators $L''_k = \sum_{n \geq 1} v_{nk}L_n$, if necessary, we can suppose that both $(L_\ell\rho^{1/2})_{\ell \geq 1}$ and $(\rho^{1/2}L_k^*)_{k \geq 1}$ are *orthogonal* bases of the same closed linear space. Note that

$$\mathrm{tr}(\rho^{1/2}(L'')^*_k\rho^{1/2}\theta(L'')^*_j\theta) = \sum_{m,n \geq 1} \bar{v}_{nk}v_{mj}\mathrm{tr}(\rho^{1/2}L_n^*\rho^{1/2}\theta L_m^*\theta)$$

and the operator R , in the new GKSL representation, transforms into V^*RV which is also self-adjoint.

Expanding $\rho^{1/2}\theta L_k^*\theta$ with respect to the orthogonal basis $(L_\ell\rho^{1/2})_{\ell \geq 1}$, for all $k \geq 1$, we have

$$\rho^{1/2}\theta L_k^*\theta = \sum_{\ell \geq 1} \frac{\mathrm{tr}(\rho^{1/2}L_\ell^*\rho^{1/2}\theta L_k^*\theta)}{\|L_\ell\rho^{1/2}\|_{HS}^2} L_\ell\rho^{1/2} \quad (23)$$

i.e. $\rho^{1/2}L_k^* = \sum_{\ell} y_{k\ell} L_{\ell} \rho^{1/2}$ with a matrix Y of complex numbers $y_{k\ell}$.

Clearly, since C is diagonal and commutes with R , we have $y_{k\ell} = (RC^{-1})_{k\ell} = (C^{-1}R)_{k\ell}$. It follows then from 3 above that Y coincides with the unitary operator $C^{-1}R$ and condition 2 of Theorem 21 holds. Moreover, Y is self-adjoint because both R and C are. $\square$

As an immediate consequence of the commutation of R and C we have the following parallel of Corollary 17 for the SQDB condition

Corollary 23 *Suppose that a QMS $\mathcal{T}$ satisfies the SQDB- θ condition. For every special GKSL representation of $\mathcal{L}$ with operators $L_{\ell}\rho^{1/2}$ orthogonal as Hilbert-Schmidt operators on $\mathfrak{h}$ if $\text{tr}(\rho^{1/2}L_{\ell}^*\rho^{1/2}\theta L_k^*\theta) \neq 0$ for a pair of indices $k, \ell \geq 1$, then $\text{tr}(\rho L_{\ell}^* L_{\ell}) = \text{tr}(\rho L_k^* L_k)$.*

When the time reversal θ is given by the conjugation $\theta u = \bar{u}$ (with respect to some orthonormal basis of $\mathfrak{h}$), $\theta x^* \theta$ is equal to the transpose x^T of x and we find the following

Corollary 24 *$\mathcal{T}$ satisfies the SQDB- θ condition if and only if there exists a special GKSL representation of $\mathcal{L}$, with operators G, L_k , such that:*

1. $\rho^{1/2}G^T = G\rho^{1/2}$;
2. $\rho^{1/2}L_k^T = \sum_j u_{kj} L_j \rho^{1/2}$ for some unitary self-adjoint $(u_{kj})_{kj}$.

6 SQDB- θ for QMS on $M_2(\mathbb{C})$

In this section, as an application, we find a standard form of a special GKSL representation of the generator $\mathcal{L}$ of a QMS on $M_2(\mathbb{C})$ satisfying the SQDB- θ . The faithful invariant state ρ , in a suitable basis, can be written in the form

$$\rho = \begin{pmatrix} \nu & 0 \\ 0 & 1 - \nu \end{pmatrix} = \frac{1}{2}(\sigma_0 + (2\nu - 1)\sigma_3), \quad 0 < \nu < 1$$

where σ_0 is the identity matrix and $\sigma_1, \sigma_2, \sigma_3$ are the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

The time reversal θ is the usual conjugation in the above basis.

In order to determine the structure of the operators G and L_k satisfying conditions of Corollary 24 we find first a convenient basis of $M_2(\mathbb{C})$. We

choose then a basis of eigenvectors of the linear map $X \rightarrow \rho^{1/2} X^T \rho^{-1/2}$ in $M_2(\mathbb{C})$ given by $\sigma_0, \sigma_1^\nu, \sigma_2^\nu, \sigma_3$ where

$$\sigma_1^\nu = \begin{pmatrix} 0 & \sqrt{2\nu} \\ \sqrt{2(1-\nu)} & 0 \end{pmatrix}, \quad \sigma_2^\nu = \begin{pmatrix} 0 & -i\sqrt{2\nu} \\ i\sqrt{2(1-\nu)} & 0 \end{pmatrix}.$$

Indeed, $\sigma_0, \sigma_1^\nu, \sigma_3$ (resp. σ_2^ν) are eigenvectors of the eigenvalue 1 (resp. -1).

Every special GKSL representation of $\mathcal{L}$ is given by (see [11] Lemma 6.1)

$$L_k = -(2\nu - 1)z_{k3}\sigma_0 + z_{k1}\sigma_1^\nu + z_{k2}\sigma_2^\nu + z_{k3}\sigma_3, \quad k \in \mathcal{J} \subseteq \{1, 2, 3\}$$

with vectors $z_k := (z_{k1}, z_{k2}, z_{k3})$ ($k \in \mathcal{J}$) linearly independent in $\mathbb{C}^3$.

The SQDB- θ holds if and only if G, L_k satisfy

$$(i) \quad G = \rho^{1/2} G^T \rho^{-1/2},$$

$$(ii) \quad L_k = \sum_{j \in \mathcal{J}} u_{kj} \rho^{1/2} L_j^T \rho^{-1/2} \text{ for some unitary self-adjoint } U = (u_{kj})_{k,j \in \mathcal{J}}.$$

Now, if $\mathcal{J} \neq \emptyset$, since every unitary self-adjoint matrix is diagonalizable and its spectrum is contained in $\{-1, 1\}$, it follows that $U = W^* D W$ for some unitary matrix $W = (w_{ij})_{i,j \in \mathcal{J}}$ and some diagonal matrix D of the form

$$\text{diag}(\epsilon_1, \dots, \epsilon_{|\mathcal{J}|}), \quad \epsilon_i \in \{-1, 1\}, \quad (24)$$

where $|\mathcal{J}|$ denotes the cardinality of $\mathcal{J}$. Therefore, replacing the L_k 's by operators $L'_k := \sum_{j \in \mathcal{J}} w_{kj} L_j$ if necessary, we can take U of the form (24).

We now analyze the structure of L_k 's corresponding to the different (diagonal) forms of U . By condition (ii) we have either $L_k = \rho^{1/2} L_k^T \rho^{-1/2}$ or $L_k = -\rho^{1/2} L_k^T \rho^{-1/2}$; an easy calculation shows that

$$L_k = \rho^{1/2} L_k^T \rho^{-1/2} \quad \text{if and only if} \quad z_{k2} = 0 \quad (25)$$

and

$$L_k = -\rho^{1/2} L_k^T \rho^{-1/2} \quad \text{if and only if} \quad z_{k1} = z_{k3} = 0. \quad (26)$$

Therefore, the linear independence of $\{z_j : j \in \mathcal{J}\}$ forces U to have at most two eigenvalues equal to 1 and at most one equal to -1 and, with a suitable choice of a phase factor for each L_k , we can write

$$L_k = (1 - 2\nu)r_k\sigma_0 + r_k\sigma_3 + \zeta_k\sigma_1^\nu \quad \text{for } k = 1, 2 \text{ and } r_k \in \mathbb{R}, \zeta_k \in \mathbb{C} \quad (27)$$

$$L_3 = r_3\sigma_2^\nu \quad r_3 \in \mathbb{R}. \quad (28)$$

Clearly L_1 and L_2 are linearly independent if and only if $r_1\zeta_2 \neq r_2\zeta_1$. This, together with non triviality conditions leaves us, up to a change of indices, with the following possibilities:

- a) $|\mathcal{J}| = 1, U = 1$ then $\mathcal{J} = \{1\}$ with $r_1\zeta_1 \neq 0$,
- b) $|\mathcal{J}| = 1, U = -1$ then $\mathcal{J} = \{3\}$ with $r_3 \neq 0$,
- c) $|\mathcal{J}| = 2, U = \text{diag}(1, 1)$ then $\mathcal{J} = \{1, 2\}$ with $r_1\zeta_1 r_2\zeta_2 \neq 0, r_1\zeta_2 \neq r_2\zeta_1$,
- d) $|\mathcal{J}| = 2, U = \text{diag}(1, -1)$ then $\mathcal{J} = \{1, 3\}$, with $r_3 \neq 0, r_1\zeta_1 \neq 0$,
- e) $|\mathcal{J}| = 3, U = \text{diag}(1, 1, -1)$ then $\mathcal{J} = \{1, 2, 3\}$ with $r_1\zeta_2 \neq r_2\zeta_1, r_3 \neq 0, r_1\zeta_1 r_2\zeta_2 \neq 0$.

To conclude, we analyze condition (i). If $G = (g_{jk})_{1 \leq j, k \leq 2}$ then statement (i) is equivalent to

$$\sqrt{\nu} g_{21} = \sqrt{1 - \nu} g_{12}. \quad (29)$$

Since $G = -iH - 2^{-1} \sum_k L_k^* L_k$ with $H = \sum_{j=1}^3 v_j \sigma_j$, $v_j \in \mathbb{R}$, and $\sum_k L_k^* L_k$ is equal to the sum of a term depending only on σ_0 and σ_3 plus

$$\sum_{k=1,2} 2r_k \begin{pmatrix} 0 & \zeta_k \sqrt{2\nu}(1 - \nu) - \bar{\zeta}_k \nu \sqrt{2(1 - \nu)} \\ \bar{\zeta}_k \sqrt{2\nu}(1 - \nu) - \zeta_k \nu \sqrt{2(1 - \nu)} & 0 \end{pmatrix}$$

in the case $\mathcal{J} \neq \emptyset$ the identity (29) holds if and only if

$$\begin{cases} v_1 (\sqrt{1 - \nu} - \sqrt{\nu}) &= -\sqrt{2\nu(1 - \nu)} (\sqrt{1 - \nu} + \sqrt{\nu})^2 \sum_{k=1}^2 r_k \Im \zeta_k \\ v_2 (\sqrt{1 - \nu} + \sqrt{\nu}) &= -\sqrt{2\nu(1 - \nu)} (\sqrt{1 - \nu} - \sqrt{\nu})^2 \sum_{k=1}^2 r_k \Re \zeta_k \end{cases}. \quad (30)$$

On the other hand, when $\mathcal{J} = \emptyset$, condition (29) is equivalent to $\sqrt{\nu}(v_1 + iv_2) = \sqrt{1 - \nu}(v_1 - iv_2)$, i.e.

$$v_1 (\sqrt{1 - \nu} - \sqrt{\nu}) = 0, \quad v_2 = 0 \quad (31)$$

Therefore we have the following possible standard forms for $\mathcal{L}$.

Theorem 25 *Let L_1, L_2, L_3 be as in (27), (28), $H = \sum_{j=1}^3 v_j \sigma_j$ with v_1, v_2 as in (30) and $v_3 \in \mathbb{R}$. The QMS $\mathcal{T}$ satisfies the SQDB- θ if and only if there exists a special GKSL representation of $\mathcal{L}$ given, up to phase factors multiplying L_1, L_2, L_3 , in one of the following ways:*

- o) H with $v_1 = v_2 = 0$ if $\nu \neq 1/2$, and $v_1 \in \mathbb{R}, v_2 = 0$ if $\nu = 1/2$,
- a) H, L_1 with $r_1\zeta_1 \neq 0$,
- b) H, L_3 with $r_3 \neq 0$,
- c) H, L_1, L_2 with $r_1\zeta_1 r_2\zeta_2 \neq 0$ and $r_1\zeta_2 \neq r_2\zeta_1$,

- d) H, L_1, L_3 with $r_3 \neq 0$ and $r_1\zeta_1 \neq 0$,
- e) H, L_1, L_2, L_3 with $r_1\zeta_2 \neq r_2\zeta_1$, $r_1\zeta_1 r_2\zeta_2 \neq 0$ and $r_3 \neq 0$.

Roughly speaking, the standard form of $\mathcal{L}$ corresponds, up to degeneracies when some of the parameter vanish or when some linear dependence arises, to the case e).

We know that a QMS satisfying the usual (i.e. with pre-scalar product with $s = 0$) QDB- θ condition must commute with the modular group. Moreover, when this happens, the SQDB- θ and QDB- θ conditions are equivalent (see e.g. [6], [11]).

We finally show how the generators of a QMSs on $M_2(\mathbb{C})$ satisfying the usual QDB- θ condition can be recovered by a special choice of the parameters $r_1, r_2, r_3, \zeta_1, \zeta_2$ in Theorem 25 describing the generator of a QMS satisfying the SQDB- θ condition.

To this end, we recall that $\mathcal{T}$ fulfills the QDB- θ when $\text{tr}(\rho x \mathcal{T}_t(y)) = \text{tr}(\rho \theta y^* \theta \mathcal{T}_t(\theta x^* \theta))$ for all $x, y \in \mathcal{B}(\mathfrak{h})$. In [11] we classified generators of QMS on $M_2(\mathbb{C})$ satisfying the QDB condition without time reversal (i.e., formally, replacing θ by the identity operator, that is, of course, not antiunitary). The same type of arguments show that, disregarding trivialisations that may occur when some of the parameters below vanish, QMSs on $M_2(\mathbb{C})$ satisfying the QDB- θ condition have the following standard form

$$\begin{aligned} \mathcal{L}(x) = & i[H, x] - \frac{|\eta|^2}{2} (L^2 x - 2LxL + xL^2) \\ & - \frac{|\lambda|^2}{2} (\sigma^- \sigma^+ x - 2\sigma^- x \sigma^+ + x \sigma^- \sigma^+) - \frac{|\mu|^2}{2} (\sigma^+ \sigma^- x - 2\sigma^+ x \sigma^- + x \sigma^+ \sigma^-), \end{aligned} \quad (32)$$

where $H = h_0 \sigma_0 + h_3 \sigma_3$ ($h_0, h_3 \in \mathbb{R}$), $L = -(2\nu - 1)\sigma_0 + \sigma_3$, $\sigma^\pm = (\sigma_1 \pm i\sigma_2)/2$ and, changing phases if necessary, λ, μ, η can be chosen as *non-negative real* numbers satisfying

$$\lambda^2(1 - \nu) = \nu\mu^2. \quad (33)$$

Choosing $r_1 = \eta, \zeta_1 = 0$ we find immediately that the operator L in (32) coincides with the operator L_1 in (27). Moreover, choosing $r_2 = 0$ we find $v_2 = 0$ and also $v_1 = 0$ for $\nu \neq 1/2$. A straightforward computation yields

$$\begin{pmatrix} \lambda \sigma_+ \\ \mu \sigma_- \end{pmatrix} = \begin{pmatrix} \lambda/(2\zeta_2\sqrt{2\nu}) & i\lambda/(2r_3\sqrt{2\nu}) \\ \mu/(2\zeta_2\sqrt{2(1-\nu)}) & -i\mu/(2r_3\sqrt{2(1-\nu)}) \end{pmatrix} \begin{pmatrix} L_2 \\ L_3 \end{pmatrix}$$

and the above 2×2 matrix is unitary if we choose $\zeta_2 = \lambda/(2\sqrt{\nu})$, $r_3 = i\mu/(2\sqrt{1-\nu}) = i\zeta_2$ because of (33) and changing the phase of r_3 in order to find a unitary that is also self-adjoint.

This shows that we can recover the standard form (32) choosing H , L_1, L_2, L_3 as in Theorem 25 e) with $r_1 = \eta, \zeta_1 = 0, r_2 = 0, \zeta_2 = \lambda/(2\sqrt{\nu}), r_3 = i\mu/(2\sqrt{1-\nu}), v_1 = v_2 = 0$.

Appendix

We denote by $\ell^2(J)$ the Hilbert space of complex-valued, square summable sequences indexed by a finite or countable set J .

Lemma 26 *Let $\mathcal{J}$ be a complex separable Hilbert space and let $(\xi_j)_{j \in J}, (\eta_j)_{j \in J}$ be two Hilbertian bases of $\mathcal{J}$ satisfying $\sum_{j \in J} \|\xi_j\|^2 < \infty, \sum_{j \in J} \|\eta_j\|^2 < \infty$. The complex matrices $A = (a_{jk})_{j,k \in J}, B = (b_{jk})_{j,k \in J}, C = (c_{jk})_{j,k \in J}$ given by*

$$a_{jk} = \langle \xi_j, \xi_k \rangle, \quad b_{jk} = \langle \xi_j, \eta_k \rangle, \quad c_{jk} = \langle \eta_j, \eta_k \rangle$$

define trace class operators on $\ell^2(J)$ satisfying $B^ A^{-1} B = C$. Moreover A and C are self-adjoint and positive.*

Proof. Note that

$$\sum_{j,k \geq 1} |b_{jk}|^2 \leq \sum_{j,k \geq 1} \|\xi_j\|^2 \cdot \|\eta_k\|^2 = \sum_j \|\xi_j\|^2 \cdot \sum_k \|\eta_k\|^2 < \infty$$

Therefore B defines a Hilbert-Schmidt operator on $\ell^2(J)$.

In a similar way A and C define Hilbert-Schmidt operators on $\ell^2(J)$ that are obviously self-adjoint. These are also positive because for any sequence $(z_m)_{m \in J}$ of complex numbers with $z_m \neq 0$ for a finite number of indices m at most we have

$$\sum_{m,n \in J} \bar{z}_m a_{mn} z_n = \sum_{m,n \in J} \bar{z}_m \langle \xi_m, \xi_n \rangle z_n = \left\| \sum_{m \in J} z_m \xi_m \right\|^2 \geq 0.$$

Moreover, they are trace class because

$$\sum_{j \in J} a_{jj} = \sum_{j \in J} \|\xi_j\|^2 < \infty, \quad \sum_{j \in J} c_{jj} = \sum_{j \in J} \|\eta_j\|^2 < \infty.$$

Finally, we show that B is also trace class. By the spectral theorem, we can find a unitary $V = (v_{kj})_{k,j \in J}$ on $\ell^2(J)$ such that $V^* A V$ is diagonal. The series $\sum_{m \in J} v_{mj} \xi_m$ is norm convergent because

$$\left\| \sum_m v_{mj} \xi_m \right\|^2 = \sum_{m,n \in J} \bar{v}_{nj} a_{nm} v_{mj} = (V^* A V)_{jj}.$$

The series $\sum_{m \in J} v_{mj} \xi_m$ is norm convergent as well for a similar reason. Therefore, putting $\xi'_j = \sum_{m \in J} v_{mj} \xi_m$ and $\eta'_j = \sum_{m \in J} v_{mj} \eta_m$ we find immediately $(V^*AV)_{kj} = \langle \xi'_k, \xi'_j \rangle = 0$ for $j \neq k$, $(V^*AV)_{jj} = \|\xi'_j\|^2$ and

$$\begin{aligned} (V^*BV)_{kj} &= \sum_{m,n} \bar{v}_{mk} v_{nj} \langle \xi_m, \eta_j \rangle = \langle \xi'_k, \eta'_j \rangle, \\ (V^*CV)_{kj} &= \sum_{m,n} \bar{v}_{mk} v_{nj} \langle \eta_m, \eta_j \rangle = \langle \eta'_k, \eta'_j \rangle. \end{aligned}$$

As a consequence, the following identity

$$\begin{aligned} (V^*B^*A^{-1}BV)_{kj} &= ((V^*B^*V)(V^*AV)^{-1}(V^*BV))_{kj} \\ &= \sum_{m \in J} (V^*B^*V)_{km} ((V^*AV)_{mm})^{-1} (V^*BV)_{mj} \\ &= \sum_{m \in J} \left\langle \eta'_k, \frac{\xi'_m}{\|\xi'_m\|} \right\rangle \left\langle \frac{\xi'_m}{\|\xi'_m\|}, \eta'_j \right\rangle \\ &= \langle \eta'_k, \eta'_j \rangle = (V^*CV)_{kj} \end{aligned}$$

holds because $(\xi'_m/\|\xi'_m\|)_{m \in J}$ is an orthonormal basis of $\mathcal{J}$.

This proves that $V^*B^*A^{-1}BV = V^*CV$ i.e. $B^*A^{-1}B = C$. It follows that $|A^{-1/2}B| = C^{1/2}$ is Hilbert-Schmidt as well as $A^{-1/2}B$ and $B = A^{1/2}(A^{-1/2}B)$ is trace class being the product of two Hilbert-Schmidt operators. $\square$

Acknowledgment. The financial support from the MIUR PRIN 2007 project “Quantum Probability and Applications to Information Theory” is gratefully acknowledged.

References

- [1] S. Albeverio and D. Goswami, A Remark on the Structure of Symmetric Quantum Dynamical Semigroups, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **5** (2002) 571–579.
- [2] L. Accardi and K. Imafuku, Dynamical detailed balance and local KMS condition for non-equilibrium states, *Internat. J. Modern Phys. B* **18** no. 4-5, 435–467 (2004).
- [3] L. Accardi and A. Mohari, Time Reflected Markov Processes, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, **2**, 397-426 (1999).

- [4] G.S. Agarwal, Open quantum Markovian systems and the microreversibility, *Z. Physik* **258** no. 5, 409–422 (1973).
- [5] R. Alicki, On the detailed balance condition for non-Hamiltonian systems, *Rep. Math. Phys.*, **10**, 249–258 (1976).
- [6] F. Cipriani, Dirichlet Forms and Markovian Semigroups on Standard Forms of von Neumann Algebras, *J. Funct. Anal.*, **147**, 259–300 (1997).
- [7] F. Cipriani, Dirichlet Forms on Noncommutative Spaces. *Quantum Potential Theory*, 161–276, Lecture Notes in Math., 1954, Springer, 2008.
- [8] E.B. Davies, J.M. Lindsay, Non-commutative symmetric Markov semigroups, *Math. Z.* **210** (1992), 379–411.
- [9] J. Dereziński, R. Fruboës, Fermi Golden Rule and Open Quantum Systems. In S. Attal et al. (eds.) *Open Quantum Systems III*, Lecture Notes in Mathematics **1882**, 67–116, Springer 2006.
- [10] F. Fagnola and V. Umanità, Detailed Balance, time reversal and generators of Quantum Markov Semigroups, *M. Zametki*, **84** (2008), no. 1, 108–116 (Russian); translation *Math. Notes* **84** (2008), no. 1–2, 108–115.
- [11] F. Fagnola and V. Umanità, Generators of Detailed Balance Quantum Markov Semigroups, *Inf. Dimens. Anal. Quantum Probab. Relat. Top.*, **10** (3), 335 - 363 (2007).
- [12] F. Fagnola and V. Umanità, On two quantum versions of the detailed balance condition. To appear in: Noncommutative harmonic analysis with applications to probability (M. Bożejko, et al. eds.), Banach Center Publications, Polish Academy of Sciences 2009.
- [13] M. Fukushima, Y. Oshima, M. Takeda, *Dirichlet Forms and Symmetric Markov Processes*, de Gruyter Studies in Mathematics 19, 1994.
- [14] S. Goldstein and J.M. Lindsay, Beurling-Deny condition for KMS-symmetric dynamical semigroups, *C.R.Acad.Sci.Paris*, **317**, 1053–1057 (1993).
- [15] S. Goldstein and J.M. Lindsay, KMS symmetric semigroups, *Math. Z.*, **219**, 591–608 (1995).
- [16] V. Gorini, A. Kossakowski and E.C.G. Sudarshan, Completely positive dynamical semigroups of N -level systems, *J. Math. Phys.* **17**, 821–825 (1976).

- [17] D. Guido, T. Isola, S. Scarlatti, Non-symmetric Dirichlet forms on semifinite von Neumann algebras, *J. Funct. Anal.* **135** (1996), no. 1, 50–75.
- [18] A. Kossakowski, A. Frigerio, V. Gorini and M. Verri, “Quantum detailed balance and KMS condition,” *Comm. Math. Phys.*, **57**, 97-110 (1977).
- [19] G. Lindblad, On the generators of Quantum Dynamical Semigroups, *Comm. Math. Phys.* **48**, 119–130 (1976).
- [20] W.A. Majewski, On the relationship between the reversibility of detailed balance conditions, *Ann. Inst. Henri Poincaré, A*, **39**, 45-54 (1983).
- [21] W.A. Majewski, The detailed balance condition in quantum statistical mechanics, *J. Math. Phys.* **25** no. 3, 614–616 (1984).
- [22] W.A. Majewski and R.F. Streater, Detailed balance and quantum dynamical maps. *J. Phys. A: Math. Gen.*, **31**, 7981-7995, (1998).
- [23] Y.M. Park, Remarks on the structure of Dirichlet forms on standard forms of von Neumann Algebras, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, **8**, 179-197 (2005).
- [24] K. R. Parthasarathy, *An introduction to quantum stochastic calculus*, *Monographs in Mathematics* 85, Birkhäuser-Verlag, Basel 1992.
- [25] D. Petz, Conditional Expectation in Quantum Probability. In L. Accardi and W. von Waldenfels (eds.) *Quantum Probability and Applications III*. Proceedings, Oberwolfach 1987. LNM **1303** p. 251–260. Springer 1988.
- [26] J.L. Sauvageot, Quantum Dirichlet forms, differential calculus and semi-groups. In L. Accardi, W. von Waldenfels (eds.), *Quantum Probability and Applications V*. LNM **1442**, 334–346, Springer 1990.
- [27] P. Talkner, The failure of the quantum regression hypothesis, *Annals of Physics*, **167** (2), 390-436 (1986).